



BRITISH COLUMBIA  
CENTRE *for* EXCELLENCE  
*in* HIV/AIDS

# HIV MONITORING QUARTERLY REPORT **FOR NORTHERN HEALTH**

SECOND QUARTER 2014



BC Centre for Disease Control  
An agency of the Provincial Health Services Authority



First Nations Health Authority  
Health through wellness



fraserhealth



Interior Health  
For your whole life



island health



northern health  
the northern way of caring



How you want to be treated.



Promoting wellness. Ensuring care.



## Foreword

As part of the BC Centre for Excellence (BC-CFE) in HIV/AIDS's mandate to evaluate the outcomes of STOP HIV/AIDS programming in BC, we have developed quarterly HIV/AIDS monitoring reports. These reports provide up-to-date data on a variety of key HIV-related surveillance and treatment indicators. Selection of these indicators was achieved through a collaborative process with various Health Authority (HA) representatives. There are six reports in total, one for each HA and one for the province of BC as a whole. In addition, there is a technical report which explains how each HIV indicator is calculated. Data used in these reports come from the British Columbia Centre for Disease Control (BCCDC), MSP billings, hospitalization data from the Discharge Abstract Database, the Sunquest Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory, Providence Health Care laboratory and the BC-CFE Drug Treatment Program (DTP) Database.

The objectives of these reports are to:

1. Provide timely HA-specific information on key HIV indicators which will guide and inform HIV leaders and innovators in the development of future HIV interventions and programs which will ultimately lead to decreasing the burden of HIV in BC. The indicators will reflect ongoing or past successful public health interventions and highlight areas in the HIV care spectrum which require further attention and support.
2. Highlight limitations in our current data due to incomplete or time lagged data and to develop future strategies to improve complete and timely data capture.

These reports are produced for the benefit of individual HA's. As such, we are enthusiastic about your involvement and cooperation regarding the development of these monitoring reports. Please forward your comments and queries to Irene Day, Director of Operations at the BC-CFE at [iday@cfenet.ubc.ca](mailto:iday@cfenet.ubc.ca).

## List of Indicators

Indicator 1. Testing Episodes

Indicator 2. HIV Testing Rate

Indicator 3. New HIV Diagnoses

Indicator 4. Stage of HIV Infection at Diagnosis

Indicator 5. HIV Cascade of Care

Indicator 6. Programmatic Compliance Score (PCS)

Indicator 7. New Antiretroviral Starts

Indicator 8. CD4 Cell Count at ART Initiation

Indicator 9. Active and Inactive Drug Treatment Program Participants

Indicator 10. Antiretroviral Adherence Level

Indicator 11. Resistance Testing Results by Resistance Category

Indicator 12. AIDS-Defining Illness

Indicator 13. HIV-Related Mortality

# Table of Contents

## Acknowledgements and Contributions

### BC Provincial STOP Program:

### A Note on Monitoring and Interpreting HIV Indicators

#### **Indicator 1      HIV Testing Episodes**

- Figure 1.1      HIV Test Episodes for Northern Health, 2009 Q3–2014 Q2
- Figure 1.2      HIV Test Episodes for Northern Health by Gender and Prenatal Status, 2009 Q3–2014 Q2
- Figure 1.3      HIV Test Episodes for Northern Health by Age Category, 2009 Q3–2014 Q2
- Figure 1.4      Point-of-Care HIV Tests for Northern Health, 2010 Q4–2014 Q2
- Figure 1.5      HIV Test Episodes by HSDA for Northern Health, 2009 Q3–2014 Q2

#### **Indicator 2      HIV Testing Rates**

- Figure 2.1      Rate of HIV Testing for Northern Health and HSDA's, 2009–2013
- Figure 2.2      Rate of HIV Testing for Northern Health by Gender, 2009–2013
- Figure 2.3      Rate of HIV Testing for Northern Health by Age Category, 2009–2013

#### **Indicator 3      New HIV Diagnoses**

- Figure 3.1      New HIV Diagnoses for Northern Health, 2009 Q3–2014 Q2
- Figure 3.2      New HIV Diagnoses for Northern Health by Gender, 2009 Q3–2014 Q2
- Figure 3.3      New HIV Diagnoses for Northern Health by Age Category, 2009 Q3–2014 Q2
- Figure 3.4      New HIV Diagnoses for Northern Health by Exposure Category, 2009 Q3–2013 Q4
- Figure 3.5      New HIV Diagnoses for Northern Health by HSDA, 2009 Q3–2014 Q2

#### **Indicator 4      Stage of HIV Infection at Diagnosis**

- Table 1      Staging Classifications of Infection at Time of HIV Diagnosis Based on CDC HIV Surveillance Case Definitions
- Figure 4.1      Stage of HIV Infection at Diagnosis for Northern Health, 2010–2013
- Figure 4.2      Stage of HIV Infection at Diagnosis for Northern Health by Gender, 2010–2013
- Figure 4.3      Stage of HIV Infection at Diagnosis for Northern Health by Age Category, 2010–2013
- Figure 4.4      Stage of HIV Infection at Diagnosis for Northern Health by Exposure Category, 2010–2013

#### **Indicator 5      HIV Cascade of Care**

- Figure 5.1      Estimated Cascade of Care for Northern Health, Year Ending 2014 Q2
- Figure 5.2      Estimated Cascade of Care for Northern Health by Gender, Year Ending 2014 Q2

Figure 5.3	Estimated Cascade of Care for Northern Health by Age Category, Year Ending 2014 Q2
Figure 5.4	Estimated Cascade of Care for Northern Health by MSM Status, Year Ending 2014 Q2
Figure 5.5	Estimated Cascade of Care for Northern Health by Age Category and MSM Status, Year Ending 2014 Q2
Figure 5.6	Estimated Cascade of Care for Northern Health by History of IDU, Year Ending 2014 Q2
Figure 5.7	Estimated Cascade of Care for Northern Health by HSDA, Year Ending 2014 Q2
<b>Indicator 6</b>	<b>Programmatic Compliance Score (PCS)</b>
Table 2	Probability of Mortality Based on the Programmatic Compliance Score
Figure 6.1	PCS Components for Northern Health, 2012 Q3–2014 Q2 <ul style="list-style-type: none"> <li>First-Year CD4 Measurement</li> <li>First-Year VL measurement</li> <li>Baseline Resistance Testing</li> <li>Recommended Antiretroviral Therapy (ART)</li> <li>Baseline CD4 <math>\geq 200</math> cells/<math>\mu</math>L</li> <li>Suppression at 9 Months</li> </ul>
Figure 6.2	Historical Trends for PCS Score for Northern Health, 2012 Q3–2014 Q2
<b>Indicator 7</b>	<b>New Antiretroviral Therapy Starts in Northern Health</b>
Figure 7	BC-CfE Drug Treatment Program Enrollment: New Antiretroviral Participants for Northern Health, 2012 Q3–2014 Q2
<b>Indicator 8</b>	<b>CD4 Cell Count at ART Initiation</b>
Figure 8	CD4 Cell Count at ART Initiation for Northern Health, 2012 Q3–2014 Q2
<b>Indicator 9</b>	<b>Active and Inactive Drug Treatment Program (DTP) Participants</b>
Table 3	Distribution of People on ART in Northern Health, 2014 Q2
Figure 9	Active and Inactive DTP Participants for Northern Health, 2012 Q3–2014 Q2
<b>Indicator 10</b>	<b>Antiretroviral Adherence</b>
Figure 10	Distribution of Individuals by Adherence Level in 1st Year of Therapy, Based on Pharmacy Refill Compliance for Northern Health, 2012 Q3–2014 Q2
<b>Indicator 11</b>	<b>Resistance Testing and Results</b>
Figure 11	Cumulative Resistance Testing Results by Resistance Category for Northern Health, 2012 Q3–2014 Q2
<b>Indicator 12</b>	<b>AIDS-Defining Illness</b>
Figure 12	AIDS Case Rate and Reports for Northern Health, 2006–2013
<b>Indicator 13</b>	<b>HIV-Related Mortality</b>
Figure 13	HIV-Related Deaths by Year for Northern Health, 2004–2011

# Acknowledgements and Contributions



BRITISH COLUMBIA  
CENTRE *for* EXCELLENCE  
*in* HIV/AIDS

**British Columbia Centre for Excellence in HIV/AIDS (BC-CFE):** The BC-CFE is responsible for the conception, preparation and ongoing review of this quarterly report. The BC-CFE provides the data and outputs for Indicators 5 (Hiv Cascade of Care), 6 (Programmatic Compliance Score), 7 (New Antiretroviral Starts), 8 (CD4 Cell Count at ART Initiation), 9 (Active and Inactive Drug Treatment Program Participants), 10 (Antiretroviral Adherence Level), 11 (Resistance Testing Results by Resistance Category), 12 (AIDS-Defining Illness), and 13 (HIV-Related Mortality). The BC-CFE database provides PVL and CD4 cell count testing data, as well as ART use. All PVL measurements in BC are performed at the St Paul's Hospital virology laboratory, thus PVL data capture is 100%. An estimated 80% of all CD4 count measurements performed in the province are captured in the BC-CFE data holdings. The STOP HIV/AIDS Technical Monitoring Committee-BC-CFE is responsible for oversight of the monitoring report. Motoi Matsukura writes and compiles the monitoring report. Guillaume Colley, Dr. Viviane Lima and Nada Gataric perform analysis of Indicators 5–13. James Nakagawa is responsible for publishing and editing. This report was conceived and guided by Dr. Julio Montaner.



BC Centre for Disease Control  
An agency of the Provincial Health Services Authority

**British Columbia Centre for Disease Control (BCCDC):** The BCCDC provides the data and outputs for Indicator 1 (Hiv Testing Episodes), Indicator 2 (Hiv Testing Rate), Indicator 3 (New Hiv Diagnoses), Indicator 4 (Stage of Hiv at Diagnosis) and Indicator 12 (AIDS-Defining Illness). The BCCDC is the single provincial agency that centralizes all HIV surveillance through the Public Health Microbiology and Reference Laboratory, which does more than 90% of all HIV screening tests in BC and all confirmatory testing. Theodora Consolacion and Dr. Mark Gilbert are responsible for outputs for Indicators 1–4.

## Other Data Sources:

The above databases were supplemented with:

- (I) The BC Vital Statistics database which was used to calculate Indicator 5. The Hiv Cascade of Care and Indicator 13. HIV-Related Mortality.
- (II) Linkage and preparation of the de-identified individual-level database used for calculating Indicator 5. The Hiv Cascade of Care was facilitated by the British Columbia Ministry of Health.
- (III) The Statistics Canada database: BC and HIV-positive population counts were acquired through the statistics Canada website to calculate HIV-specific mortality rates for Indicator 13. HIV-Related Mortality.

# Membership of the STOP HIV/AIDS Technical Monitoring Committee–BC-CfE

Dr. Rolando Barrios, *Chair*, BC-CfE

Kate Heath, BC-CfE

Bohdan Nosyk, BC-CfE

Viviane Dias Lima, BC-CfE

Irene Day, BC-CfE

Dr. Mark Gilbert, BCCDC

Dr. Mel Kradjen, BCCDC

Stephanie Konrad, FHA

Joanne Nelson, FNHA

Jennifer May-Hadford, IHA

James Haggerstone, NHA

Dr. Neora Pick, PHSA

Dr. Reka Gustafson, VCHA

Melanie Rusch, VIHA



# The Seek and Treat for Optimal Prevention (STOP) HIV/AIDS BC Provincial Program: A Note on Monitoring and Interpreting HIV Indicators

The Seek and Treat for Optimal Prevention (STOP) of HIV/AIDS programme is a provincial initiative to improve HIV diagnosis and care delivery in BC through increased HIV-specific funding to all HSDA's across BC. The STOP provincial programme is an expansion of a four-year STOP pilot project which was implemented in two Health Service Delivery Areas in March 2010; the Vancouver HSDA which bears the largest burden of the HIV epidemic in the province and the Northern Interior HSDA which bears a high burden of HIV-related mortality. The STOP pilot project demonstrated the urgent need for improved efforts in early diagnosis of HIV and timely initiation of antiretroviral therapy (ART) initiation.

The expansion to a province-wide programme was announced on November 30th 2013 by the BC Ministry of Health with roll out of funding beginning on April 1st, 2013. This funding is intended to be used in the implementation and evaluation of HIV-related diagnosis and care initiatives within individual HA's. Goals of the project include: 1. A reduction in the number of new HIV infections in BC; 2. Improvements in the quality, effectiveness, and reach of HIV prevention services; 3. An increase in early diagnosis of HIV; 4. A reduction in AIDS cases and HIV-related mortality.

The goals of HA-led STOP-funded initiatives are to work toward achieving these goals. To these ends some outcome measures or indicators of progress have been drafted that should be considered in the design and implementation phases of these initiatives.

# HIV Testing Episodes and Rates


In this section, the number of HIV test episodes and point of care (POC) HIV tests conducted each quarter in BC is shown. In general terms the goal is to increase the number of tests performed and to maximize testing efficiency. Test episodes are allocated by region according to where the test is performed.

## Indicator 1. HIV Testing Episodes

Figure 1.1 HIV Test Episodes for Northern Health, 2009 Q3–2014 Q2



Figure 1.2 HIV Test Episodes by Gender and Prenatal Status for Northern Health, 2009 Q3–2014 Q2<sup>1</sup>



<sup>1</sup> NB: Testing does not include point of care tests.

Figure 1.3 HIV Test Episodes by Age Category for Northern Health, 2009 Q3–2014 Q2<sup>1,2</sup>





Figure 1.4 Point-of-Care HIV Tests for Northern Health, 2010 Q4–2014 Q2




<sup>2</sup> Data Source: The BC Public Health Microbiology and Reference Laboratory (BCPHMRL) courtesy of the BC Centre for Disease Control (BCCDC).

**Limitations:**

- 1 Repeat tests in individuals who test using various identifiers may not be identified and these individuals may be counted more than once.
- 2 POC testing data is available from the fourth quarter of 2010 and onwards.

Figure 1.5 HIV Test Episodes for Northern Health by HSDA, 2009 Q3–2014 Q2





## Indicator 2. HIV Testing Rates

Figure 2.1 Rate of HIV Testing for Northern Health and HSDAs, 2009–2013 <sup>1</sup>




Figure 2.2 Rate of HIV Testing by Gender for Northern Health, 2009–2013 <sup>1</sup>





Figure 2.3 Rate of HIV Testing by Age Category for Northern Health, 2009–2013 <sup>1</sup>



<sup>1</sup> NB: Testing does not include point of care tests.

# New HIV Diagnoses

Trends in HIV diagnoses by gender and exposure category are described. Interpreting HIV diagnoses must be done with consideration that trends are influenced by both changes in testing rate as well as changes in transmission rates. It is important to note that new HIV diagnoses cases and rates are not synonymous with HIV incidence as a person may have become infected with HIV long before they tested positive for HIV. However, as there is no reliable method for measuring HIV incidence we follow trends in HIV diagnoses.

## Indicator 3. New HIV Diagnoses

Figure 3.1 New HIV Diagnoses for Northern Health, 2009 Q3–2014 Q2 <sup>3</sup>





Figure 3.2 New HIV Diagnoses for Northern Health by Gender, 2009 Q3–2014 Q2



<sup>3</sup> Data Source: BCCDC “By Provider Address” is graphed as dashed line in same colour.



Figure 3.3 New HIV Diagnoses for Northern Health by Age Category, 2009 Q3–2014 Q2<sup>3</sup>




Figure 3.4 New HIV Diagnoses for Northern Health by Exposure Category, 2009 Q3–2013 Q4<sup>3,4</sup>





Figure 3.5 New HIV Diagnoses for Northern Health by HSDA, 2009 Q3–2014 Q2<sup>3</sup>



<sup>3</sup> Data Source: BCCDC. "By Provider Address" is graphed as dashed line in same colour.

<sup>4</sup> MSM=men who have sex with men; IDU= injection drug user; HET=heterosexual. NIR=No identified risk/exposure.

# Stage of HIV infection at diagnosis

Classification of stage of HIV infection, in the absence of information regarding recent testing history, is reliant on clinical information available at the time of diagnosis, including first CD4+ cell count, laboratory results suggestive of acute HIV infection, and clinical presentation with an AIDS-defining illness (Table 1). The benefits of Treatment as Prevention (TasP) are maximized when antiretroviral therapy (ART) is initiated at high CD4 cell counts. Accordingly, it is preferable that individuals newly diagnosed with HIV be in the early stages of HIV infection (stage 0 or 1) to allow for early ART initiation.

*N.B. Interpretation of stage of HIV infection at diagnosis should proceed with caution. Early increases in diagnosis at late stage (i.e., low CD4 counts) may represent a “catching up” of previously missed long term infected individuals rather than a trend toward diagnosis at later stage of infection.*

## Indicator 4. Stage of HIV Infection at Diagnosis

Table 1 Staging Classifications of Infection at Time of HIV Diagnosis Based on CDC HIV Surveillance Case Definitions

Stage	Criteria		
0	Laboratory criteria met for acute HIV infection, or previous negative or indeterminate HIV test within 180 days of first confirmed positive HIV test.		
1	Stage 0 not met	CD4 ≥500	and No AIDS case report
2a		CD4 350–499	
2b		CD4 200–349	
3		( CD4 <200	or AIDS case report )
Unknown		No available CD4	and No AIDS case report

Figure 4.1 Stage of HIV Infection at Diagnosis for Northern Health, 2010–2013 <sup>5</sup>




Figure 4.2 Stage of HIV Infection at Diagnosis by Gender for Northern Health, 2010–2013 <sup>5</sup>




Figure 4.3 Stage of HIV Infection at Diagnosis by Age Category for Northern Health, 2010–2013 <sup>5</sup>





Figure 4.4 Stage of HIV Infection at Diagnosis by Exposure Category for Northern Health, 2010–2013 <sup>5,6</sup>



6 MSM=men who have sex with men; IDU= injection drug user; HET=heterosexual. NIR=No identified risk/exposure.

## Indicator 5. HIV Cascade of Care

The success of seek, test, treat and retain (STTR) strategies like STOP is reliant on early diagnosis of HIV, linking newly diagnosed HIV-positive persons with ongoing care, retaining persons in HIV-care; initiating ART based on best evidenced practices and maintaining optimal ART adherence to ensure a suppressed viral load. These stages of HIV-care can be summarized as: 1. HIV diagnosis, 2. Linkage to HIV care, 3. Retention in HIV care, 4. On ART and 5. Achieving a suppressed VL; collectively, they are referred to as the cascade of care. Leakage between any of these stages of HIV-care means a reduction in the potential of ART as a benefit to the HIV-positive individual and as an HIV transmission prevention method on a population level. Thus, when interpreting trends in the cascade of care, we strive to see increases along each step of the cascade of care (ie. reduced attrition) with the ultimate goal being 100% within each stage of the cascade. Monitoring the Cascade of Care provides a picture as to where deficiencies lie in the delivery and uptake of HIV-care. In this section we present the cascade of care for the year 2012 in BC overall and stratified by sex and age for each Health Authority.

Figure 5.1 Estimated Cascade of Care for Northern Health, Year Ending 2014 Q2 <sup>7</sup>





Figure 5.2 Estimated Cascade of Care for Northern Health by Gender, Year Ending 2014 Q2 <sup>8</sup>



<sup>7,8</sup> Data is for the period 2013 Q3–2014 Q2.

Data Sources:

<sup>1</sup> British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).


<sup>2</sup> Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider.

If the most recent HA of residence is not updated then the designated HA may be incorrect.


NB: Transgender has been assigned to their biological sex.

Figure 5.3      Estimated Cascade of Care for Northern Health by Age Category, Year Ending 2014 Q2 <sup>9</sup>



<sup>9</sup> Data is for the period 2013 Q3–2014 Q2.  
Data Sources:  
1 British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).  
2 Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).  
Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider.  
If the most recent HA of residence is not updated then the designated HA may be incorrect.

Figure 5.4 Estimated Cascade of Care for Northern Health by MSM Status, Year Ending 2014 Q2 <sup>10</sup>




<sup>10</sup> Data is for the period 2013 Q3–2014 Q2.

Data Sources:

- 1 British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- 2 Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider. If the most recent HA of residence is not updated then the designated HA may be incorrect.

Figure 5.5 Estimated Cascade of Care for Northern Health by Age Category and MSM Status, Year Ending 2014 Q2 <sup>11</sup>




<sup>11</sup> Data is for the period 2013 Q3–2014 Q2.

Data Sources:

- 1 British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- 2 Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider. If the most recent HA of residence is not updated then the designated HA may be incorrect.

Figure 5.6 Estimated Cascade of Care for Northern Health by History of IDU, Year Ending 2014 Q2 <sup>12</sup>



<sup>12</sup> Data is for the period 2013 Q3–2014 Q2.


Data Sources:

- 1 British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- 2 Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider. If the most recent HA of residence is not updated then the designated HA may be incorrect.



Figure 5.7 Estimated Cascade of Care for Northern Health by HSDA, Year Ending 2014 Q2 <sup>13</sup>



<sup>13</sup> Data is for the period 2013 Q3–2014 Q2.

Data Sources:

- 1 British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- 2 Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider. If the most recent HA of residence is not updated then the designated HA may be incorrect.

## Indicator 6. The Programmatic Compliance Score (PCS)

The Programmatic Compliance Score (PCS) is a summary measure of risk of future death, immunologic failure and virologic failure from all causes for people who are starting ART for the first time. It is composed of patient- and physician-driven effects. PCS scores range from 0–6 with higher scores indicative of poorer health outcomes and greater risk of death. Table 1 provides mortality, immunologic failure and virologic failure probabilities for given PCS scores. We interpret an individual with a  $PCS \geq 4$  as being 22 times more likely to die, almost 10 times more likely to have immunologic failure and nearly 4 times as likely to demonstrate virologic failure compared to those individuals with a PCS score of 0. A detailed description of how the PCS score is calculated and its validation can be found in the technical report. In short, PCS scores are calculated by summing the results (yes=1, no=0) of six un-weighted non-performance indicators based on IAS–USA treatment guidelines:

1. having <3 CD4 cell count tests in the first year after starting antiretroviral therapy (ART);
2. having <3 plasma viral load (VL) tests in the first year after starting ART;
3. not having drug resistance testing done prior to starting ART;
4. starting on a non-recommended ART regimen;
5. starting therapy with  $CD4 < 200$  cells/ $\mu$ L; and
6. not achieving viral suppression within 9 months since ART initiation.

In this section we provide PCS scores and their components over time for the province of BC. A decline to 0%, (**i.e., all individuals having a score of 0**) is the eventual goal.

Table 2. The Probability of Mortality, Immunologic Failure and Virologic Failure based on the Programmatic Compliance Score

Programmatic Compliance Score	Mortality Risk Ratio (95% Confidence Interval)	Immunologic Failure Risk Ratio (95% CI)	Virologic Failure Risk Ratio (95% CI)
0 (Best score)	1 (–)	1 (–)	1 (–)
1	3.81 (1.73–8.42)	1.39 (1.04–1.85)	1.32 (1.05–1.67)
2	7.97 (3.70–17.18)	2.17 (1.54–3.04)	1.86 (1.46–2.38)
3	11.51 (5.28–25.08)	2.93 (1.89–4.54)	2.98 (2.16–4.11)
4 or more (Worst score)	22.37 (10.46–47.84)	9.71 (5.72–16.47)	3.80 (2.52–5.73)

Reference: Lima VD, Le A, Nosyk B, Barrios R, Yip B, et al. (2012) Development and Validation of a Composite Programmatic Assessment Tool for HIV Therapy. *PLoS ONE* 7(11): e47859. doi:10.1371/journal.pone.0047859

Figure 6.1 PCS Components for Northern Health, 2012 Q3–2014 Q2 <sup>14</sup>





Figure 6.2 Historical Trends for PCS Score for Northern Health, 2012 Q3–2014 Q2 <sup>14,15</sup>



<sup>14</sup> Data Source: British Columbia Centre for Excellence Drug Treatment Program (DTP) Database. Limitations: CD4 cell count capture is approximately 80%.


<sup>15</sup> Each quarter's data is calculated as the sum of the 4 quarters leading up to it. e.g. 2013 Q1 is calculated from 2012 Q2 – 2013 Q1. NB: A score of 0 is the best score and a score of 4 or more is the worst score.

## Antiretroviral Uptake

In this section we present trends in ART uptake, the number and proportion of new HIV treatment initiations and the number of active and inactive DTP participants. Trends in ART uptake should be interpreted under the consideration of changing BC HIV treatment guidelines. BC HIV treatment guidelines are updated regularly by the BC-CfE Therapeutic Guidelines Committee and reflect those of the International AIDS Society. Most recent changes were made in 2012 and HIV treatment is now recommended for all HIV-positive adults regardless of CD4 cell count; as evidence demonstrates that early initiation of HIV treatment maximizes both the individual's health outcomes as well as the potential of ART as a form of HIV transmission prevention at a population level. As such, trends in the number and proportion of persons on ART and new ART starts (in both naïve and experienced persons) are expected to increase over time at higher CD4 cell counts.

### Indicator 7. New Antiretroviral Therapy Starts in Northern Health

Figure 7 BC-CfE Drug Treatment Program Enrollment: New ART Participants in Northern Health, 2012 Q3–2014 Q2<sup>16</sup>



### Indicator 8. CD4 Cell Count at ART Initiation

Figure 8 CD4 Cell Count at ART Initiation of ART-Naïve DTP Participants in Northern Health, 2012 Q1–2013 Q4<sup>17</sup>

The majority of cells in this figure have  $n \leq 5$ , which is considered statistically insignificant as well as a possible risk to patient privacy. For this reason, this figure has been omitted. Authorized parties may contact the British Columbia Centre for Excellence in HIV/AIDS to obtain this information.

<sup>16</sup> Data Source: Drug Treatment Program Database

Limitation: DTP participants are designated to an HA based on most current residence provided by the participant.

<sup>17</sup> Data Source: Drug Treatment Program Database


Limitations: CD4 cell count data is approximately 80% complete.

## Indicator 9. Active and Inactive DTP Participants

Table 3. Distribution of People on ART for Northern Health, 2014 Q2 <sup>18</sup>

Age	< 30	13
	30–39	48
	40–49	68
	≥ 50	90
Gender	Male	134
	Female	85
Exposure	MSM	26
	IDU	129
Total		219

Figure 9 Active and Inactive DTP Participants in Northern Health, 2012 Q3–2014 Q2 <sup>19</sup>



<sup>18</sup> Data Source: Drug Treatment Program Database

Limitation: DTP participants are designated to an HA based on most current residence provided by the participant.

Definitions:

'On antiretroviral therapy' defined as being on treatment in the current quarter

'Unknown/not stated' defined as being on treatment in the current quarter, and city of residence unknown

<sup>19</sup> Active DTP participants: are those who are prescribed one or more drugs in the last six months.

Inactive DTP Participants: Persons no longer prescribed drugs through the HIV/AIDS Drug Treatment Program in the last quarter.

## Antiretroviral Adherence Level

In this section we present trends in prescription refill adherence levels for individuals in their first year of treatment. Given that the benefits of ART are compromised in the presence of imperfect ART adherence, we expect to see the proportion of persons on ART achieving *near perfect adherence* (ie.  $\geq 95\%$ ) to increase with time. Furthermore, it is important that trends in the proportion of ART users achieving prescription refill adherence of  $\geq 95\%$  keep pace with new ART starts and increase among those continuing on ART.

### Indicator 10. Antiretroviral Adherence

Figure 10      Distribution of Individuals by Adherence Level in 1st  
Year of Therapy, Based on Pharmacy Refill Compliance  
for Northern Health, 2012 Q3–2014 Q2 <sup>20</sup>


The majority of cells in this figure have  $n \leq 5$ , which is considered statistically insignificant as well as a possible risk to patient privacy. For this reason, this figure has been omitted. Authorized parties may contact the British Columbia Centre for Excellence in HIV/AIDS to obtain this information.

---

<sup>20</sup>    *Data Source: Drug Treatment Program Database*  
     *Limitation: Prescription refill adherence is used as a proxy for patient adherence.*

# Indicator 11. Resistance Testing and Results

In this section, we present trends in cumulative resistance testing by resistance category: **Suppressed** (where a DTP participant's viral load is too low to be genotyped); **Wild Type** (where no HIV treatment resistances were discovered), **Never Genotyped**, and Resistances to **one**, **two** or **three** HIV treatment classes. Resistance testing prior to ART initiation is recommended in the BC HIV treatment primary care guidelines. Thus, it is expected that trends over time should find all persons enrolled in the DTP to have been genotyped. Trends over time should also show an increase in the proportion of DTP participants achieving a suppressed status and an increase in resistance testing should not lead to an increase in the number of ART resistances occurring.




<sup>21</sup> Data Source: Drug Treatment Program Database

Limitation: DTP participants are designated to an HA based on most current residence provided by the participant.

## Indicator 12. AIDS-Defining Illness

Improvements in ART and the expansion of ART province-wide has led to very low numbers of recorded AIDS cases across BC. However, interpreting trends in AIDS cases is challenging as AIDS reporting is passive in BC and it is likely that they are under reported across all Health Authorities. In addition to under reporting, methods of reporting AIDS cases are inconsistent across HA's and do not truly reflect the current reality of new AIDS diagnoses. Efforts will need to be made to improve under and inconsistent reporting of AIDS cases across all HA's. The table below shows AIDS cases using three definitions. First, AIDS cases were defined as the number of physician-reported AIDS defining illness (ADI) in a given year. AIDS case reporting is a passive process and physicians can voluntarily report AIDS cases to the BCCDC or DTP. As such, we have plotted both **BCCDC reports** and **DTP reported AIDS cases**. We also show the proportion of persons **initiating ART with a CD4<200 cells/μL**.

Figure 12 AIDS Case Rate and Reports for Northern Health, 2006–2013<sup>22</sup>




<sup>22</sup> Data Source: DTP AIDS cases are obtained from the Drug Treatment Program Database; BCCDC AIDS cases are obtained from the BC-CDC; CD4<200 at ART initiation data came from the DTP database.

*Limitation:* AIDS case reporting was investigated using 3 definitions: First, using AIDS cases reported in AIDS case report forms from the DTP; Second, using AIDS cases reported via the BCCDC and third, using a CD4 cell count of <200 cells/μL at time of ART initiation using DTP data. AIDS case reporting is passive in BC, thus; AIDS case reporting is not well captured. The DTP sends out AIDS reporting forms to physicians annually. The BCCDC uses DTP AIDS case reports as well as physician AIDS case reports made directly to the BCCDC. Interpreting AIDS case reports should be done with these limitations in mind. AIDS data is updated annually as very few AIDS cases reports are reported in general and trends would be difficult to notice if reported quarterly.



### Indicator 13. HIV-Related Mortality

Evidence indicates that individuals who initiate treatment with recommended ART in a timely fashion may live near normal lifespans. Excess mortality among HIV positive persons is, therefore, an important measure of HIV care with a goal of minimizing HIV-related mortality in British Columbia.



<sup>23</sup> Data Source: BC Vital Statistics

*Limitation:*

1. DTP participants are designated to an HA based on most current residence provided by the participant.
2. Mortality data is updated annually.
3. The most recent available data was used.

# Appendices

Indicator 1: <b>Test Episodes (thousands)</b>		2009		2010		2011		2012		2013		2014	
		Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2
Northern Health		2.7	2.5	2.9	2.6	2.7	2.5	3.0	2.7	2.9	2.8	3.4	3.1
Gender	Female	1.8	1.8	1.9	1.8	1.8	1.7	2.0	1.8	1.9	1.9	2.2	2.0
	Male	0.8	0.7	0.9	0.8	0.8	0.7	0.9	0.8	0.8	0.8	1.1	1.0
	Other	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female (Prenatal)		0.9	1.0	0.9	0.8	0.9	0.8	1.0	0.9	1.0	0.9	1.0	0.9
Female (Non-prenatal)		0.9	0.9	1.0	1.0	0.9	0.9	1.0	0.9	1.0	1.2	1.1	1.2
Age	< 30	1.4	1.3	1.4	1.3	1.4	1.2	1.4	1.3	1.4	1.4	1.6	1.5
	30–39	0.7	0.7	0.8	0.7	0.7	0.7	0.8	0.7	0.8	0.7	0.9	0.8
	40–49	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.4
	≥ 50	0.2	0.2	0.4	0.3	0.3	0.2	0.4	0.3	0.3	0.3	0.5	0.4
POC HIV Tests (not in thousands)							45	19	49	54	52	83	74
Northeast		0.7	0.6	0.7	0.7	0.6	0.6	0.7	0.7	0.8	0.7	0.9	0.8
Northern Interior		1.3	1.2	1.5	1.3	1.3	1.3	1.5	1.4	1.4	1.5	1.8	1.6
Northwest		0.7	0.7	0.8	0.7	0.7	0.6	0.7	0.6	0.7	0.6	0.8	0.7

Indicator 2: **Rate of HIV Testing per 100,000**

		2009	2010	2011	2012	2013
All Northern Health		3515.1	3435.5	3583.1	4089.0	4391.3
Northeast		3793.4	3522.7	3797.9	4011.6	4017.1
Northern Interior		3482.5	3420.2	3612.2	4248.5	4719.9
Northwest		3324.2	3384.3	3326.9	3856.3	4121.5
Gender	Female	4925.8	4790.9	4991.1	5528.6	5833.4
	Male	2086.5	2085.6	2190.8	2664.2	2964.7
Age	< 30	4321.4	4202.9	4402.9	4840.9	4888.4
	30–39	7555.5	7329.5	7697.9	8064.0	8291.6
	40–49	2591.3	2666.6	2801.6	3294.2	3707.2
	≥ 50	1169.3	1202.0	1264.2	1929.2	2625.0

Indicator 3: <b>New HIV Diagnoses</b>		2009		2010		2011		2012		2013		2014	
		Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2
Northern Health	By Client Residence	5	7	4	3	7	2	4	10	5	5	5	4
	By Provider Address	5	8	4	3	6	2	4	9	5	6	5	4
Gender	Female	3	3	2	1	2	0	1	5	1	3	1	2
	Male	2	4	2	2	5	2	3	5	4	2	4	2
Age	< 30	1	3	0	1	0	0	0	4	1	2	1	0
	30–39	1	1	2	1	2	0	1	4	1	2	0	1
	40–49	2	2	1	0	3	2	2	1	2	0	1	2
	≥ 50	1	1	1	1	2	0	1	1	1	3	1	1
Exposure	MSM	0	0	0	0	1	0	0	1	1	1	0	0
	IDU	4	4	4	1	2	2	3	5	1	3	3	2
	HET	1	2	0	2	3	0	1	3	3	1	2	2
	Other	0	0	0	0	0	0	0	0	0	0	0	0
	NIR/Unknown	0	1	0	0	1	0	0	1	0	0	0	1
Northeast	By Client Residence	0	0	0	2	0	0	1	1	0	0	0	0
	By Provider Address	0	0	0	2	0	0	1	1	0	0	0	0
Northern Interior	By Client Residence	2	6	1	0	5	2	0	5	4	4	2	4
	By Provider Address	3	7	1	0	4	2	0	4	4	4	2	4
Northwest	By Client Residence	3	1	3	1	2	0	3	4	1	1	3	0
	By Provider Address	2	1	3	1	2	0	3	4	1	2	3	0

**Indicator 4: Stage of HIV Infection at Baseline**

	Northern Health				Female				Male				< 30 years				30–39 years				40–49 years			
	'10	'11	'12	'13	'10	'11	'12	'13	'10	'11	'12	'13	'10	'11	'12	'13	'10	'11	'12	'13	'10	'11	'12	'13
Stage 0	2	3	0	2	0	2	0	1	2	1	0	1	0	1	0	0	0	1	0	1	1	1	0	0
Stage 1	2	5	5	3	0	2	1	0	2	3	4	3	0	1	1	0	0	1	0	3	2	1	1	0
Stage 2a	4	2	2	1	1	1	1	0	3	1	1	1	0	2	0	1	4	0	0	0	0	0	0	0
Stage 2b	1	6	2	3	1	2	1	2	0	4	1	1	1	0	0	0	0	3	1	1	0	2	1	0
Stage 3	6	4	2	7	4	1	0	1	2	3	2	6	0	1	0	0	1	2	0	2	3	1	1	1
Unknown	0	4	1	1	0	2	0	0	0	2	1	1	0	1	0	0	0	2	0	0	0	0	1	1
Total	15	24	12	17	6	10	3	4	9	14	9	13	1	6	1	1	5	9	1	7	6	5	4	2

	≥ 50 years				MSM				IDU				Heterosexual				Other Exposure				NIR/Unknown			
	'10	'11	'12	'13	'10	'11	'12	'13	'10	'11	'12	'13	'10	'11	'12	'13	'10	'11	'12	'13	'10	'11	'12	'13
Stage 0	1	0	0	1	0	0	0	1	1	2	0	0	0	1	0	1	0	0	0	0	1	0	0	0
Stage 1	0	2	3	0	0	0	1	1	2	4	3	1	0	1	1	0	0	0	0	1	0	0	0	0
Stage 2a	0	0	2	0	1	0	0	0	2	0	1	1	1	2	1	0	0	0	0	0	0	0	0	0
Stage 2b	0	1	0	2	0	2	0	0	0	0	1	1	1	3	1	1	0	1	0	1	0	0	0	0
Stage 3	2	0	1	4	0	1	0	1	4	2	1	0	2	1	1	6	0	0	0	0	0	0	0	0
Unknown	0	1	0	0	0	0	0	0	0	3	0	1	0	0	0	0	0	0	0	0	0	1	1	0
Total	3	4	6	7	1	3	1	3	9	11	6	4	4	8	4	8	0	1	0	2	1	1	1	0

**Indicator 5: HIV Cascade of Care**

		DIAGNOSED	LINKED	RETAINED	ON ART	ADHERENT	SUPPRESSED
Northern Health		270	252	230	211	186	114
Age Category	< 30	20	14	13	12	10	7
	30–39	56	53	48	43	38	19
	40–49	80	77	70	64	54	30
	≥ 50	115	108	98	92	84	58
Age Category and MSM Status	MSM	< 30	≤ 5	≤ 5	≤ 5	≤ 5	≤ 5
		30–39	≤ 5	≤ 5	≤ 5	≤ 5	≤ 5
		40–49	11	11	9	8	5
		≥ 50	16	15	14	13	8
	Non-MSM	< 30	9	8	7	4	3
		30–39	38	36	33	31	15
		40–49	46	45	41	34	19
		≥ 50	67	66	60	56	38
	Unknown	< 30	10	5	5	5	3
		30–39	16	15	10	6	4
		40–49	22	21	14	12	6
		≥ 50	31	27	18	15	12
Gender	Male	165	152	138	126	115	69
	Female	105	99	92	85	71	45
Injection Drug Use	IDU	143	139	137	129	114	68
	Non-IDU	82	78	67	62	54	36
	Unknown	46	34	26	20	18	10
MSM Status	MSM	30	29	27	25	23	14
	Non-MSM	160	155	150	140	125	75
	Unknown	80	68	53	46	38	25
Health Authority	Northeast	33	30	19	19	16	8
	Northern Interior	172	160	152	142	123	73
	Northwest	66	61	58	50	47	33

**Indicator 6: Programmatic Compliance Score (PCS)**

	2012 Q3	Q4	2013 Q1	Q2	Q3	Q4	2014 Q1	Q2
< 3 CD4 Tests	25.0%	30.0%	32.0%	34.6%	29.2%	18.8%	12.5%	6.3%
< 3 Viral Load Tests	25.0%	30.0%	24.0%	30.8%	33.3%	25.0%	25.0%	12.5%
No Baseline Genotype	2.8%	0.0%	4.0%	3.8%	4.2%	6.3%	0.0%	0.0%
Baseline CD4 < 200 cells/μL	38.9%	36.7%	24.0%	23.1%	20.8%	18.8%	37.5%	37.5%
Non-Recommended ART	5.6%	6.7%	8.0%	7.7%	8.3%	12.5%	6.3%	6.3%
Non Viral suppression at 9 Mo.	63.9%	70.0%	76.0%	76.9%	75.0%	62.5%	62.5%	50.0%
PCS Score: 0	6	4	2	3	3	3	3	5
PCS Score: 1	14	12	12	10	9	7	7	6
PCS Score: 2	6	4	4	5	5	3	3	3
PCS Score: 3	8	8	6	6	6	2	2	2
PCS Score: 4 or more	2	2	1	2	1	1	1	0
<b>Total (n=)</b>	<b>36</b>	<b>30</b>	<b>25</b>	<b>26</b>	<b>24</b>	<b>16</b>	<b>16</b>	<b>16</b>

**Indicator 7: New DTP ARV Participants**

First Starts	7	2	4	4	8	7	5	2
Experienced Starts	5	8	6	9	8	5	4	7

**Indicator 8: CD4 Cell Count at ART Initiation for ARV-Naïve DTP Participants**

CD4 ≥ 500	2	–	–	–	3	1	–	–
CD4 350–499	2	–	–	–	0	1	–	–
CD4 200–349	1	–	–	–	2	2	–	–
CD4 50–199	0	–	–	–	2	3	–	–
CD4 < 50	2	–	–	–	1	0	–	–
<i>CD4 Median (cells/μL)</i>	<i>370</i>	<i>–</i>	<i>–</i>	<i>–</i>	<i>250</i>	<i>270</i>	<i>–</i>	<i>–</i>
<b>Total (n=)</b>	<b>7</b>	<b>≤ 5</b>	<b>≤ 5</b>	<b>≤ 5</b>	<b>8</b>	<b>7</b>	<b>≤ 5</b>	<b>≤ 5</b>

**Indicator 9: Active and Inactive DTP Participants**

Active DTP Participants	181	184	185	190	201	208	212	219
Inactive DTP Participants	46	44	48	44	36	38	39	36

**Indicator 10: Antiretroviral Adherence**

≥ 95%	4	–	–	–	6	–	–	–
80% to < 95%	4	–	–	–	0	–	–	–
40% to < 80%	1	–	–	–	0	–	–	–
< 40%	0	–	–	–	0	–	–	–
<b>Total (n=)</b>	<b>9</b>	<b>≤ 5</b>	<b>≤ 5</b>	<b>≤ 5</b>	<b>6</b>	<b>≤ 5</b>	<b>≤ 5</b>	<b>≤ 5</b>

**Indicator 11: Resistance Testing and Results**

Suppressed	81	91	107	97	93	93	124	97
Wild Type	37	35	41	34	36	33	30	31
Never Genotyped	3	1	2	1	3	2	1	0
1-Class	12	13	6	9	10	8	10	7
2-Class	1	1	3	1	4	3	1	2
3-Class	0	0	0	0	0	0	0	0
<b>Total (n=)</b>	<b>134</b>	<b>141</b>	<b>159</b>	<b>142</b>	<b>146</b>	<b>139</b>	<b>166</b>	<b>137</b>

**Indicator 12: AIDS-Defining Illness**

	2006	2007	2008	2009	2010	2011	2012	2013
CD4 < 200 at Cases	12	15	14	16	10	13	7	10
ART initiation <i>Rate per 100,000</i>	<i>4.3</i>	<i>5.3</i>	<i>4.9</i>	<i>5.6</i>	<i>3.5</i>	<i>4.5</i>	<i>2.4</i>	<i>3.4</i>
AIDS Cases Cases	≤ 5	≤ 5	≤ 5	6	6	≤ 5	≤ 5	≤ 5
(DTP Reports) <i>Rate per 100,000</i>	<i>0.0</i>	<i>1.8</i>	<i>1.1</i>	<i>2.1</i>	<i>2.1</i>	<i>1.7</i>	<i>1.0</i>	<i>1.7</i>
AIDS Cases Cases	≤ 5	≤ 5	≤ 5	8	≤ 5	6	≤ 5	–
(BCCDC Reports) <i>Rate per 100,000</i>	<i>1.4</i>	<i>1.8</i>	<i>1.1</i>	<i>2.8</i>	<i>0.7</i>	<i>2.1</i>	<i>0.3</i>	<i>–</i>

**Indicator 13: HIV-Related Mortality**

	2004	2005	2006	2007	2008	2009	2010	2011
Northern Health	7	6	7	8	6	≤ 5	≤ 5	≤ 5
Per 100 HIV+ Population	2.28	1.92	2.21	2.47	1.81	0.89	1.45	1.42
Per 100,000 Population	2.30	1.95	2.42	2.75	2.11	1.05	1.73	1.72