

STOP HIV/AIDS Pilot Project

QUARTERLY INDICATORS REPORT: 1 April through 30 June 2012 (Q2)

SUBMITTED TO:
The BC Ministry of Health

SUBMITTED BY:
Dr. Mark Gilbert and Dr. Kate Heath
On behalf of The STOP HIV/AIDS Technical (Indicators) Group

CONTRIBUTORS:
Teddy Consolacion, Sung Lee, Monika Lindegger, Stanley Wong, Amanda Yu, Nada Gataric, and Benita Yip

VERSION DATE:
July 20, 2012



BRITISH COLUMBIA
CENTRE for EXCELLENCE
in HIV/AIDS



BC Centre for Disease Control
An agency of the Provincial Health Services Authority



How you want to be treated.



Contact Information

Irene Day

STOP HIV/AIDS Leadership Committee Co-Chair

Director of Operations, BC Centre for Excellence in HIV/AIDS

(604) 806-8202

iday@cfenet.ubc.ca

613-1081 Burrard Street

Vancouver, BC, V6Z 1Y6

Reka Gustafson

STOP HIV/AIDS Leadership Committee Co-Chair

Medical Health Officer and Medical Director of Communicable Disease Control

Vancouver Coastal Health

(604) 675-3925

reka.gustafson@vch.ca

800-601 West Broadway

Vancouver, BC, V5Z 4C2

Table of Contents

Introduction	4
Indicator 1: Number of HIV tests	6
Indicator 2: Population HIV testing rate.....	9
Indicator 3: Number of new HIV diagnoses.....	12
Indicator 4: Rate of new AIDS case reports	15
Indicator 5: Percentage positivity among persons tested for HIV	17
Indicator 6a: Proportion of individuals tested for syphilis who are tested for HIV at the same clinical encounter .	19
Indicator 6b: Proportion of individuals with a new STI diagnosis who are tested for HIV within three months of STI diagnosis	22
Indicator 7: Proportion of individuals with a new HCV diagnosis who are tested for HIV within three months of HCV diagnosis	25
Indicator 9: Proportion of individuals with a new HIV diagnosis with advanced HIV disease	28
Indicator 11: Proportion of individuals with a new HIV diagnosis with acute HIV infection.....	31
Indicator 14: Proportion of individuals starting antiretroviral therapy (ART) late in the course of HIV disease	34
Indicator 18: Proportion of individuals with a new HIV diagnosis who are tested for syphilis within three months of HIV diagnosis.....	36
Indicator 21: Percentage of HIV-infected individuals who are tested for genotypic antiretroviral drug resistance prior to starting antiretroviral therapy (ART).....	39
Indicator 22: Percentage of individuals starting antiretroviral therapy (ART) who achieve HIV plasma viral load (pVL) of <200 copies/mL within nine months of therapy initiation.....	41
Indicator 23: Percentage of individuals who initiated antiretroviral therapy (ART) with a recommended therapy regimen (among those with no drug resistance)	44
Indicator 24: Percentage of individuals on antiretroviral therapy (ART) that achieve annual prescription refill adherence of >95%	47
Indicator 25: Number of physicians initiating therapy or providing HIV-related care to patients on antiretroviral therapy (ART).....	49
Indicator 26: Percentage of individuals on antiretroviral therapy (ART) who experience a serious adverse drug reaction (ADR)	51
Indicator 28: Incidence of resistance to any retroviral drug	54
Indicator 29: Proportion of individuals on antiretroviral therapy who change antiretroviral drug treatment	56
Data Tables	58
Appendix A: List of Indicators	98

Introduction

STOP HIV/AIDS Overview

The Seek and Treat for Optimal Prevention of HIV AIDS Pilot Project is a five year initiative being conducted in the Vancouver and Northern Interior Health Service Delivery Areas (HSDA) to improve access to care and treatment for all eligible HIV positive individuals living in British Columbia. The long term objective is to reduce HIV transmission risk and lower HIV incidence in B.C.

This initiative is needed because although highly active antiretroviral therapy (HAART) improves clinical outcomes, reduces transmission risk and dramatically improves life expectancy among people living with HIV/AIDS, access to HAART is suboptimal. Many individuals only access treatment late in disease and many die without ever accessing therapy at all. For some individuals failure to access treatment is a function of social, economic, cultural or medical challenges while others are simply unaware that they are HIV positive.

Stop HIV/AIDS has five specific goals: 1) To reduce the number of new HIV/AIDS diagnoses in the Vancouver and Northern Interior HSDA; 2) To improve the effectiveness of HIV screening and early detection; 3) To ensure timely access and retention to high-quality and safe HIV/AIDS care and treatment; 4) To improve the patient experience in every step of the HIV/AIDS continuum and; 5) To demonstrate system cost optimization.

In order to determine to what extent the goals of STOP are being achieved an ongoing evaluation is being conducted. This evaluation will eventually be based on monitoring 36 surveillance and clinical "indicator variables" or measures selected by the Indicators Working Group and approved by the STOP HIV/AIDS Leadership Committee. Current evaluations do not include all indicators as multiple data sources are required (Appendix A provides a complete list of indicators however, currently for a complete list of indicators). Each indicator is associated with a target level based on current trends, local or international benchmarks and/or best practices and clinical guidelines.

Caution

The progress of the STOP HIV/AIDS Pilot Project as measured by select indicators is an imperative component of project monitoring and feedback to stakeholders. In this quarterly report we describe changes in 19 key indicators in the months since the introduction of the STOP HIV/AIDS Pilot Program. In total, through data linkages some 36 indicators will eventually be included in the evaluation process.

The information provided here is correct and complete to the best of current knowledge, standards and capabilities, however, it is based on administrative, clinical, surveillance and programmatic databases which have inherent limitations. The data contained in these

databases were not originally collected for the purpose for which they are now being used and limitations arise directly from their originally intended purpose. Therefore, while each database is rich in information for select utilizations, these data should comprise only one component of our efforts to inform service delivery and policy decision-making.

We are continuing to refine our analyses to improve the quality of these indicators. This may result in small changes in the magnitude of these indicators between subsequent reports. These changes to methods are documented and accompany each indicator.

It is important to proceed with caution when interpreting trends over the short term as they are presented here. Some indicators exhibit considerable variation from one reporting period to the next. This is particularly true of estimates made for the Northern Interior Health Service Delivery Area where statistics may be based on extremely small numbers allowing for particular instability in estimates. Only by review of longer-term temporal trends (including consideration of pre-pilot fluctuations) can a complete evaluation of the direction, stability and possible future progress of each trend be evaluated. It is also important to acknowledge the inherent difficulty in ascribing changes in indicators directly to the STOP HIV/AIDS initiative given the complex, rapidly progressing nature of HIV-related care, research and service delivery in the context of a dynamic health care and data-collection systems.

Despite inherent limitations of currently available data, continual refinement of indicators and reporting strategies in conjunction with planned assessment of other data sources, integration of existing extensive datasets, and triangulation of variables will be used to construct a robust scientific platform. In this context, the observation and analyses of long-term trends will provide a powerful, complete, and accurate evaluation of the STOP HIV/AIDS Pilot Project.

Indicator 1: Number of HIV test episodes

Target:

Increase by 50%

Actual:

VAN: 22,512 testing episodes in 2012 Q2

NI: 1,583 testing episodes in 2012 Q2

Figure 1.1 Number of HIV test episodes by HSDA




Figure 1.2 Number of HIV test episodes by HSDA – Males




Figure 1.3 Number of HIV test episodes by HSDA – Females





Figure 1.4 Number of POC HIV tests by HSDA



Indicator 1 Number of HIV test episodes

Interpretations & Comments	Overall, the total number of HIV test episodes per quarter has increased in all HSDA's. HIV test episodes have decreased slightly for both males and females in Q2; the number of HIV test episodes per quarter is higher in females compared to males. The number of POC HIV tests per quarter decreased in Vancouver HSDA in Q2 and remains steady in Northern Interior HSDA.
Description of Measure	The number of HIV test episodes ordered, which is a measure of the volume of HIV tests performed in an HSDA. Data includes i) prenatal HIV tests, and ii) point of care HIV tests (delivered by STOP HIV/AIDS partner agencies).
Significance	Number of HIV test episodes ordered is a direct reflection of project initiatives related to HIV screening and may equate to increased case-finding and reduced number of individuals unaware of their HIV status. Target (50% increase, based on average 2009 Q1 to Q4) by end of STOP HIV/AIDS pilot project: Vancouver HSDA 20,932 test episodes, Northern Interior HSDA 2,013 test episodes.
Data Source(s)	<ul style="list-style-type: none"> • Misys Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory (PHSA). • Point of care HIV testing volumes from STOP HIV/AIDS partner agencies (starting in 2010 Q1). • Providence Health Care laboratory data (starting in September 2011)
Calculation Method	<ul style="list-style-type: none"> • Total number of HIV tests grouped by test episodes. A test episode consists of all HIV tests conducted for an individual in a 30-day period (as follow-up or simultaneous HIV tests may be required to clarify test results within this period). • Allocation by HSDA is based on address of ordering clinician or clinic, or if unknown, address of individual undergoing HIV testing. • Unit of analysis is number of HIV test episodes per quarter.
Limitations	Includes data for ~95% of all screening and all confirmatory HIV testing in BC. Does not include data for screening HIV tests conducted at Victoria General Hospital.
Notes	<ul style="list-style-type: none"> • POC HIV test data in Figure 1.4 are included in Figure 1.1 but not in Figures 1.2 and 1.3. • The number of POC HIV tests delivered in 2011 Q1 was affected by a recall of test kits during this period.
Revisions	<ul style="list-style-type: none"> • Number of point of care HIV tests delivered by partner agencies through STOP HIV/AIDS included. (Oct 2010) • Breakdown by gender included. (Oct 2010) • Improvement to the method for data analysis has revised the values of this indicator slightly from the November 10, 2010 report. (Jan 2011) • Inclusion of Figure 1.4 (Number of POC HIV tests by HSDA). (Jan 2011) • Inclusion of Providence Health Care laboratory data (September 2011)

Indicator 2: Population HIV testing rates

Target:	Increase by 50%	
Actual:	VAN: 6,965.0 per 100,000 in 2011	NI: 3,641.4 per 100,000 in 2011

Figure 2.1 Population HIV testing rate by HSDA




Figure 2.2 Population HIV testing rate by HSDA – Males





Figure 2.3 Population HIV testing rate by HSDA – Females



Indicator 2 Population HIV testing rates

Interpretations & Comments	In 2011, the population HIV testing rate in Vancouver HSDA has increased above historical trends; Northern Interior HSDA and other HSDAs remain within historic trends. Similar trends are observed for both males and females; the HIV testing rate is higher in females compared to males.
Description of Measure	Annual population rate of unique individuals tested for HIV.
Significance	Number of individuals tested for HIV is a direct reflection of project initiatives related to HIV screening and may equate to increased case-finding and reduced number of individuals unaware of their HIV status. Target (50% increase, based on 2009 rate) by end of STOP HIV/AIDS pilot project: Vancouver HSDA 9,722 persons tested per 100,000 population, Northern Interior HSDA 5,264 persons tested per 100,000 population.
Data Source(s)	Misys Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory (PHSA).
Calculation Method	<ul style="list-style-type: none"> • Probabilistic matching of identifiers is conducted to identify individuals having greater than one HIV test in the same year. • Denominator: Population of HSDA • Numerator: Number of unique individuals tested for HIV • Allocation by HSDA is based on address of individual undergoing HIV testing, or if unknown, address of ordering clinician or clinic. • Unit of analysis is rate of individuals tested for HIV per 100,000 population per year.
Limitations	<ul style="list-style-type: none"> • As per Indicator 1. Repeat tests in individuals who test under different identifiers (e.g., initials, pseudonyms, non-nominally) may not be identified and these individuals may be counted more than once. • Indicator #2 does not include POC test volumes or Providence Health Care laboratory data and therefore cannot be compared to trends in Indicator #1 • This indicator is limited to annual reporting as if examined on a quarterly basis one does not see a big difference from the number of HIV test episodes per quarter (as repeat HIV testing is unlikely within smaller time periods).
Notes	Would be difficult to include POC HIV test data and data from other labs in this analysis, as this would require full sharing of identifying in order to link to testing done at the Provincial Public Health Microbiology and Reference Laboratory and identify unique individuals. Total number of HIV test episodes (Indicator 1) may be preferable.
Revisions	<ul style="list-style-type: none"> • Breakdown by gender included. (Oct 2010) • Improvement to the method for data analysis has revised the values of this indicator slightly from the November 10, 2010 report. (Jan 2011)

Indicator 3: Number of new HIV diagnoses

Target: Increase during first two years then decrease

Actual: VAN: 32 persons in 2012 Q2 (by Residence) NI: 3 persons in 2012 Q2 (by Residence)

Figure 3.1 Number of new HIV diagnoses by HSDA – Allocated by RESIDENCE




Figure 3.2 Number of new HIV diagnoses by HSDA – Allocated by ORDERING CLINICIAN




Figure 3.3 Number of new HIV diagnoses by gender, BC





Figure 3.4 Number of new HIV diagnoses by Aboriginal status, BC



Indicator 3 Number of new HIV diagnoses

Interpretations & Comments	Allocation by Residence: The number of new HIV diagnoses per quarter in Vancouver HSDA decreased in 2012 Q2 and remained steady in Northern Interior HSDA. In other HSDAs, the number of new diagnoses has decreased. In Q2 there was a decrease in the number of new diagnosis in BC for males yet females remained steady. The number of new HIV diagnoses in Aboriginal people is stable.
Description of Measure	Number of individuals identified with a new diagnosis of HIV (i.e., a new positive HIV test).
Significance	The number of individuals identified with a new HIV diagnosis may be influenced by initiatives to expand HIV screening (resulting in increased case-finding and an increase in new diagnoses - may be observed during initial implementation of screening initiatives) and decreases in HIV incidence as a result of expanded HAART which would result in a decrease in new HIV diagnoses.
Data Source(s)	Provincial HIV/AIDS surveillance database at BCCDC.
Calculation Method	<ul style="list-style-type: none"> On receipt of a positive HIV test result, history of previous HIV testing is elicited from provincial databases or during public health follow-up. An individual identified with a new positive HIV test in BC is included (individuals with a previous positive HIV test inside or outside BC are excluded).¹ Allocation by HSDA is done two ways: <ul style="list-style-type: none"> Figure 3.1: by Residence - based on address of individual with new HIV diagnosis, or if unknown, address of ordering clinician or clinic. Figure 3.2: by Ordering Clinician – based on address of ordering clinician or clinic, or if unknown, by address of individual with new HIV diagnosis Unit of analysis is number of new diagnoses of HIV per quarter.
Limitations	<ul style="list-style-type: none"> This indicator is not a measure of HIV incidence (number of newly acquired HIV infections) within each time period, as an individual can be diagnosed with HIV at varying lengths of time after acquiring infection (months to years). May be difficult to interpret trends given influence of both HIV testing trends and HIV incidence on this variable. In Northern Interior HSDA, there will be greater variability for this indicator due to small numbers making trends more difficult to interpret. Ethnicity is elicited during public health follow up and there is an expected reporting delay of 6 months
Notes	<ul style="list-style-type: none"> In comparing indicator reports, the number of new HIV diagnoses for the most recent quarters may decrease. This is an expected finding as during public health follow-up, individuals identified as a new HIV diagnosis are found to have previously tested positive (e.g., in another province). The number of new HIV diagnoses allocated by Ordering Physician may more accurately represent new HIV diagnoses that occur through HIV testing services within each region (e.g., residents of FHA who test and are diagnosed through VCH services are allocated to VCH HSDA).
Revisions	<ul style="list-style-type: none"> Breakdown by gender included. (Oct 2010) Addition of allocation to HSDA by Ordering Clinician (Fig 3.2). (Jan 2011) Breakdown by Aboriginal status included. (June 2011)

¹ For HIV case definition, refer to *Annual Surveillance Report: HIV and Sexually Transmitted Infections 2008, BCCDC (Technical Appendix)*.

Indicator 4: Rate of new AIDS case reports

Target: Decrease

Actual: VAN: 3.1 per 100,000 in 2010

NI: 0.0 per 100,000 in 2010

Figure 4.1 Rate of new AIDS case reports by HSDA





Figure 4.2 Rate of new AIDS case reports by gender, BC



Indicator 4 Rate of new AIDS case reports

Interpretations & Comments	In 2010, the rate of new AIDS case reports decreased in all HSDA's. The rate of new AIDS case reports in 2009 decreased for males and remained steady in females.
Description of Measure	The rate of individuals with an AIDS case report, which indicates the first diagnosis of an AIDS defining illness in an individual with HIV infection.
Significance	Presentation with an AIDS defining illness may indicate delayed diagnosis of HIV, delays in initiation of HAART or sub-optimal management of HAART.
Data Source(s)	<ul style="list-style-type: none"> Provincial HIV/AIDS surveillance database at BCCDC. The majority of AIDS case reports are reported by the Drug Treatment Program (DTP) at the BC CfE, which submits data twice yearly to BCCDC.
Calculation Method	<ul style="list-style-type: none"> Multiple AIDS case report forms may be submitted for the same individual; only the first case report form is included in the rate of new AIDS case reports.² Denominator: Population of HSDA Numerator: Number of individuals with an AIDS case report Allocation by HSDA is based on address of the individual with an AIDS case report at the time of reporting, or if unknown, address of clinician or clinic completing the AIDS case report form. Unit of analysis is the rate of new AIDS case reports per 100,000 population per year.
Limitations	<ul style="list-style-type: none"> In BC, AIDS surveillance is based on passive reporting initiated by care providers, and under-reporting is likely. There is an expected reporting delay of up to 12 months and this indicator will only be generated at the end of the following calendar year (i.e., data for 2010 will be available in January 2012). In Northern Interior HSDA, there will be greater variability for this indicator due to small numbers making trends more difficult to interpret.
Notes	In 2010, the BCCfE as part of routine program activities received historic data on cancer-related outcomes from the BC Cancer Agency for DTP participants. New AIDS diagnoses for DTP participants occurring in the past were identified and reported to BCCDC. The number of new AIDS case reports per year has increased slightly from previous reports as a result.
Revisions	<ul style="list-style-type: none"> Breakdown by gender included. (Oct 2010)

² For AIDS case definition, refer to *Annual Surveillance Report: HIV and Sexually Transmitted Infections 2008*, BCCDC (Technical Appendix).

Indicator 5: Percentage positivity among persons tested for HIV

Target	Increase from 0.4 to 0.8 percent	
Actual	VAN: 0.26% in 20112 Q2	NI: 0.40% in 2012 Q2

Figure 5.1 Percentage positivity among persons tested for HIV by HSDA





Figure 5.2 Percentage positivity among persons tested for HIV, by gender, BC



Indicator 5 Percentage positivity among persons tested for HIV

Interpretations & Comments	In 2012 Q2 the percentage positivity decreased slightly in Vancouver HSDA Due to small numbers the trend in Northern Interior HSDA remains variable. The percentage positivity among males and females tested for HIV in 2012 Q2 decreased.
Description of Measure	The percentage of unique individuals who are tested for HIV who have a positive HIV test.
Significance	Percentage positivity may be a better reflection of the effectiveness of HIV screening and case-finding than overall test volume or new diagnoses of HIV. This indicator is influenced by HIV screening initiatives (percentage positivity may increase or decrease depending on the overall test volume and reach into populations with undiagnosed HIV infection) and decreases in HIV incidence, which would result in decreased percentage positivity.
Data Source(s)	<ul style="list-style-type: none"> • Misys Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory (PHSA). • Provincial HIV/AIDS surveillance database at BCCDC.
Calculation Method	<ul style="list-style-type: none"> • Denominator: Number of unique individuals tested for HIV • Numerator: Number of unique individuals tested for HIV who have a first positive HIV test • Allocation by HSDA is based on address of ordering clinician or clinic, or if unknown, address of individual undergoing HIV testing. • Unit of analysis is the percentage positivity of all HIV tests per quarter.
Limitations	<ul style="list-style-type: none"> • As per Indicators 1 and 2. • The numerator includes individuals who have a first positive HIV test in HIV laboratory data (repeat positive tests are excluded). Individuals having a previous positive HIV test outside of BC, or who test using different identifiers, are included in the numerator. • May be difficult to interpret significance of trends given influence of both HIV testing trends and HIV incidence on this variable. In Northern Interior HSDA, there will be greater variability for this indicator due to small numbers making trends more difficult to interpret.
Notes	
Revisions	<ul style="list-style-type: none"> • Breakdown by gender included. (Oct 2010) • Improvement to the method for data analysis has revised the values of this indicator slightly from the November 10, 2010 report. (Jan 2011)

Indicator 6a: Proportion of individuals tested for syphilis who are tested for HIV at the same clinical encounter

Target:	Increase	
Actual:	VAN: 80.4% in 2012 Q2	NI: 85.5% in 2012 Q2

Figure 6a.1 Proportion of individuals tested for syphilis who are tested for HIV at the same clinical encounter by HSDA




Figure 6a.2 Proportion of individuals tested for syphilis who are tested for HIV at the same clinical encounter by HSDA – Males





Figure 6a.3 Proportion of individuals tested for syphilis who are tested for HIV at the same clinical encounter by HSDA – Females



Indicator 6a Proportion of individuals tested for syphilis who are tested for HIV at the same clinical encounter

Interpretations & Comments	Since 2010 Q1, the proportion of individuals tested per quarter for syphilis who are tested for HIV at the same clinical encounter has been relatively stable for all regions, and overall for males and females. The magnitude of this proportion is higher for females compared to males.
Description of Measure	The percentage of individuals who are tested for syphilis who are also tested for HIV at the same clinical visit or encounter. This indicator also includes women who are undergoing prenatal testing for syphilis and HIV.
Significance	A syphilis test may indicate that an individual has risk behaviors which may also be associated with an increased risk of HIV. Ensuring all individuals getting a syphilis test are tested for HIV may lead to increased case-finding and reduce the number of individuals who are unaware of their HIV status. This may be a focus of communications with clinicians conducting HIV testing.
Data Source(s)	<ul style="list-style-type: none"> • Misys Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory (PHSA). • Provincial HIV/AIDS surveillance database at BCCDC.
Calculation Method	<ul style="list-style-type: none"> • Denominator: Number of individuals having a syphilis screening test (i.e., RPR test) • Numerator: Number of individuals having a syphilis screening test who are also testing for HIV within 14 days before or after the syphilis specimen collection date • Individuals who have previously tested positive for HIV more than 14 days before the syphilis specimen collection date are excluded from the analysis. • Allocation by HSDA is based on address of ordering clinician or clinic, or if unknown, address of individual undergoing syphilis testing. • Unit of analysis is the percentage of individuals tested for syphilis who have not previously tested positive for HIV and are tested for HIV at the same clinical encounter, by quarter.
Limitations	<ul style="list-style-type: none"> • Individuals who test for HIV using different identifiers (e.g., initials, pseudonyms, non-nominally) than are used for testing for syphilis will not be included in the numerator. • POC HIV test data and HIV test data from another laboratory not included. • For these reasons, this indicator is an under-estimate
Notes	Analysis for this indicator can only be done for syphilis testing. While looking at the proportion of individuals tested for gonorrhea or chlamydia who are also tested for HIV at the same clinical encounter would be ideal, the majority of these tests are done at private labs thus testing data is not available for analysis.
Revisions	<ul style="list-style-type: none"> • Indicator debuted. (Oct 2010) • Improvement to the method for data analysis has revised the values of this indicator slightly from the November 10, 2010 report. (Jan 2011)

Indicator 6b: Proportion of individuals with a new STI diagnosis who are tested for HIV within three months of STI diagnosis

Target:	Increase	
Actual:	VAN: 39.1% in 2012 Q1	NI: 22.7% in 2012 Q1

Figure 6b.1 Proportion of individuals with a new STI diagnosis who are tested for HIV within 3 months of STI diagnosis




Figure 6b.2 Proportion of individuals with a new STI diagnosis who are tested for HIV within 3 months of STI diagnosis – Males





Figure 6b.3 Proportion of individuals with a new STI diagnosis who are tested for HIV within 3 months of STI diagnosis – Females



Indicator 6b Proportion of individuals with a new STI diagnosis who are tested for HIV within three months of STI diagnosis

Interpretations & Comments	Since 2010 Q1, the proportion of individuals with a new STI diagnosis tested for HIV within three months of STI diagnosis has remained steady in Vancouver HSDA and Other HSDA's, and increased in Northern Interior HSDA. Similar trends are observed for females and males. The magnitude of this proportion is higher for males compared to females.
Description of Measure	The percentage of individuals with a new diagnosis of a sexually transmitted infection (STI) who are tested for HIV within 3 months of their STI diagnosis.
Significance	An STI diagnosis indicates that an individual may have risk behaviors which may also be associated with an increased risk of HIV. Recommending individuals with a new STI diagnosis are tested for HIV may lead to increased case-finding and reduce the number of individuals who are unaware of their HIV status. This may be a focus of communications with clinicians conducting HIV testing.
Data Source(s)	<ul style="list-style-type: none"> Provincial STI surveillance system at BCCDC. Misys Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory (PHSA).
Calculation Method	<ul style="list-style-type: none"> An individual with a new diagnosis of an STI is defined as an individual with a new case report for chlamydia or gonorrhea (repeat diagnoses within one month excluded). The individual's new case report for an STI will be linked to an HIV test, where available. HIV test history is identified through a probabilistic match of identifiers for STI case reports and identifiers for HIV testers. Individuals with a new STI case report who are linked to an earlier positive HIV test result are excluded from the analysis. Denominator: Number of new case reports for an STI Numerator: Number of new case reports for an STI who have a documented HIV test within 14 days before or 3 months after the date of STI diagnosis Allocation by HSDA is based on address of new case report for an STI, or if unknown, address of ordering clinician or clinic. Unit of analysis is the percentage of new case reports for an STI diagnosis who are tested within 3 months for HIV, by quarter.
Limitations	<ul style="list-style-type: none"> Individuals who test for HIV using different identifiers (e.g., initials, pseudonyms, non-nominally) than are used for an STI diagnosis will not be included in the numerator. This indicator (in comparison to 6a) involves two distinct databases and the potential for misclassification is higher. POC HIV test data and HIV test data from another laboratory are not included in the data linkage. Due to data quality limitations (which are greater than for indicator 6a) this indicator underestimates the true proportion
Notes	<ul style="list-style-type: none"> This indicator depends on linkage between two databases using probabilistic matching and may be more likely to be an underestimate (i.e., compared to Indicators 6a or 7, which are primarily a linkage within the same database).
Revisions	<ul style="list-style-type: none"> Indicator debuted. (Apr 2011)

Indicator 7: Proportion of individuals with a new HCV diagnosis who are tested for HIV within three months of HCV diagnosis

Target:	Increase	
Actual:	VAN: 54.5 % in 2011 Q3 & 4	NI: 57.4% in 2011 Q3 & 4

Figure 7.1 Proportion of individuals with a new HCV diagnosis who are tested for HIV within three months of HCV diagnosis by HSDA




Figure 7.2 Proportion of individuals with a new HCV diagnosis who are tested for HIV within three months of HCV diagnosis by HSDA – Males





Figure 7.3 Proportion of individuals with a new HCV diagnosis who are tested for HIV within three months of HCV diagnosis by HSDA – Females



Indicator 7 Proportion of individuals with a new HCV diagnosis who are tested for HIV within 3 months of HCV diagnosis

Interpretations & Comments	In 2011 Q3&4, the proportion of individuals with a new HCV diagnosis tested for HIV within three months of HCV diagnosis was stable in Vancouver HSDA and Other HSDAs, and variable in Northern Interior HSDA. Similar trends were observed overall for males and females. The magnitude of this proportion is similar for males and females.
Description of Measure	The percentage of individuals with a new diagnosis of HCV who are tested for HIV within 3 months of their HCV diagnosis.
Significance	Previous BC research on HCV and HIV co-infected persons demonstrated that most individuals were infected with HCV prior to HIV. As the majority of new HCV diagnoses are considered to be related to injection drug use, this indicator may reflect HIV testing initiatives in the IDU population.
Data Source(s)	<ul style="list-style-type: none"> • Misys Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory (PHSA). • Legacy Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory (PHSA) – used to identify previous HCV diagnoses before 2006 • Provincial HIV/AIDS surveillance database at BCCDC.
Calculation Method	<ul style="list-style-type: none"> • An individual with a new HCV diagnosis is defined as an individual with a new case report for HCV. • Denominator: Number of unique individuals with a new diagnosis of HCV. • Numerator: Number of unique individuals with a new diagnosis of HCV who have an HIV test within 14 days before or 3 months after the date of HCV diagnosis • Individuals who tested positive for HIV more than 14 days before the date of HCV diagnosis are excluded from the analysis. • Allocation by HSDA is based on address of clinician or clinic ordering HCV test, or if unknown, address of individual with new HCV diagnosis. • Unit of analysis is the percentage of individuals with a new HCV diagnosis who have not previously tested positive for HIV and are tested for HIV within 3 months, per six months.
Limitations	<ul style="list-style-type: none"> • In Northern Interior HSDA, there will be greater variability for this indicator due to small numbers making trends more difficult to interpret. • Use of partial or differing identifiers may affect linkage to HIV test results • POC HIV test data and HIV test data from other laboratories not included. • Due to data quality limitations this indicator underestimates the true proportion.
Notes	May be better indicator than Indicator 6 as have large number of HCV diagnoses, and strong validity as marker for injection drug use, which is a priority population for HIV testing through STOP HIV/AIDS.
Revisions	<ul style="list-style-type: none"> • Individuals with a previous positive HIV test excluded from analysis. (Oct 2010) • Breakdown by gender included. (Oct 2010) • Allocation by HSDA has changed from the previous report where allocation was based first on address of individual with new HCV diagnosis. To more accurately reflect testing done within each HSDA, allocation is now based first on address of clinician or clinic ordering HCV. (Oct 2010) • Access to Legacy Laboratory data has permitted identification of individuals having a HCV diagnosis prior to 2006, who are now excluded from the analysis (i.e., as not a new diagnosis). While overall trends are similar, the absolute number of new HCV diagnoses and proportion tested for HIV within three months of diagnosis per quarter are lower than the November 2010 Indicator report. (Jan 2011)

Indicator 9: Proportion of individuals with a new HIV diagnosis with advanced HIV disease

Target:	Decrease	
Actual:	VAN: 8.1% in 2010	NI: 0.0% in 2010

Figure 9.1 Proportion of individuals with a new HIV diagnosis with advanced HIV disease by HSDA




Figure 9.2 Proportion of individuals with a new HIV diagnosis with advanced HIV disease by gender, BC





Figure 9.3 Proportion of individuals with a new HIV diagnosis with advanced HIV disease by Aboriginal status, BC



Indicator 9 Proportion of individuals with a new HIV diagnosis with advanced HIV disease

Interpretations & Comments	In 2010, the proportion of individuals with a new HIV diagnosis with advanced HIV disease (AHD) was relatively stable and within the historic range in Vancouver HSDA and Other HSDAs, and more variable for Northern Interior. Among males, this proportion decreased slightly in 2010 and remained variable for females.
Description of Measure	The percentage of individuals testing newly positive for HIV who are at an advanced stage of HIV infection at the time of their HIV diagnosis.
Significance	Indicates the proportion of individuals with a new positive HIV test who test at an advanced stage of infection (i.e., diagnosis occurs years later than the time of HIV infection). These individuals have had persistent undiagnosed HIV infection which impacts on clinical care and may contribute to ongoing HIV transmission. Delays in diagnosis may be due to lack of awareness regarding risk of HIV or barriers to accessing HIV testing (i.e., HIV stigma).
Data Source(s)	Provincial HIV/AIDS surveillance database at BCCDC.
Calculation Method	<ul style="list-style-type: none"> • Probabilistic matching of identifiers is used to link AIDS and HIV case report forms. AHD at diagnosis is defined as an individual with a new diagnosis of HIV and with a linked AIDS case report form before or up to 12 months after the date of HIV diagnosis. • Denominator: Number of individuals newly diagnosed with HIV (Indicator 3) • Numerator: Number of individuals newly diagnosed with HIV and with AHD • Allocation by HSDA is based on address of individual with new HIV diagnosis, or if unknown, address of ordering clinician or clinic. • Unit of analysis is proportion of newly diagnosed individuals with AHD per year.
Limitations	<ul style="list-style-type: none"> • As per Indicator 4. There is an expected reporting delay of up to 12 months and this indicator will only be generated at the end of the following calendar year (i.e., data for 2010 will be available in January 2012). • Individuals with different identifiers on HIV and AIDS case report forms will not be identified (and are not included in the numerator). • In Northern Interior HSDA, there will be greater variability for this indicator due to small numbers making trends more difficult to interpret
Notes	<ul style="list-style-type: none"> • This indicator can be improved by consideration of first viral load and CD4+ count, which will allow for greater identification of AHD (e.g., expand AHD case definition to include all individuals with a first CD4+ count of < 200 cells/mm³). This will be achieved through data linkage with BCCFE data and is captured in Indicator 10. • In 2010, the BCCfE as part of routine program activities received historic data on cancer-related outcomes from the BC Cancer Agency for DTP participants. New AIDS diagnoses for DTP participants occurring in the past were identified and reported to BCCDC. The number of new AIDS case reports per year has increased slightly from previous reports as a result.
Revisions	<ul style="list-style-type: none"> • Breakdown by gender included. (Oct 2010) • Breakdown by Aboriginal status included. (June 2011)

AHD = advanced HIV disease

Indicator 11: Proportion of individuals with a new HIV diagnosis with acute HIV infection

Target:	Increase	
Actual:	VAN: 10 % in 2011	NI: 0% in 2011

Figure 11.1 Proportion of individuals with a new HIV diagnosis with acute HIV infection by HSDA




Figure 11.2 Proportion of individuals with a new HIV diagnosis with acute HIV infection by gender, BC





Figure 11.3 Proportion of individuals with a new HIV diagnosis with acute HIV infection by Aboriginal status, BC



Indicator 11 Proportion of individuals with a new HIV diagnosis with acute HIV infection

Interpretations & Comments	In 2011, the proportion of individuals with a new HIV diagnosis with acute HIV infection remained steady in all HSDA's. This proportion declined for females and remained steady for males. The proportion for Aboriginal persons decreased in 2011.
Description of Measure	The percentage of individuals testing newly positive for HIV who are identified as having acute HIV infection (i.e., tested up to 6-8 weeks after infection with HIV).
Significance	Individuals may test for HIV during the period of acute infection due to sero-conversion symptoms, as a result of enhanced case-finding (e.g., testing of contacts of a new index HIV case), by testing after a recent risk exposure or event, or by chance (e.g., a routine tester who tests while acutely infected). Increases in this indicator may reflect overall earlier diagnosis of HIV or increased HIV testing frequency in individuals at risk of HIV infection.
Data Source(s)	Provincial HIV/AIDS surveillance database at BCCDC.
Calculation Method	<ul style="list-style-type: none"> Acute HIV infection is defined on the basis of characteristic laboratory findings and the absence of an AIDS case report before or up to 12 months after HIV diagnosis. Denominator: All unique individuals with a new HIV diagnosis. Numerator: Number of unique individuals with a new HIV diagnosis and with acute HIV infection. Allocation by HSDA is based on address of individual with new HIV diagnosis, or if unknown, address of ordering clinician or clinic. Unit of analysis is proportion of newly diagnosed individuals with acute HIV infection per year.
Limitations	<ul style="list-style-type: none"> Ability to identify acute HIV infection depends on test window periods, which vary by type of test used (which may vary by HSDA and over time). Pooled NAAT testing is available at select clinics with gay male clients in Vancouver and contributes to increased detection of acute HIV infection in men testing at those sites. A future switch from 3rd generation to 4th generation EIA testing at the Provincial Public Health Microbiology and Reference Laboratory may influence trends. In Northern Interior HSDA, there will be greater variability for this indicator due to small numbers making trends more difficult to interpret.
Notes	
Revisions	<ul style="list-style-type: none"> Breakdown by gender included. (Oct 2010) Breakdown by Aboriginal status included. (June 2011)

Indicator 14: Proportion of individuals starting antiretroviral therapy (ART) late in the course of HIV disease

Target:	Decrease	
Actual:	VAN: 18.18%	NI: 0.00%

Figure 14.1 Proportion of individuals starting antiretroviral therapy (ART) late in the course of HIV disease by HSDA





Figure 14.2 Proportion of individuals starting antiretroviral therapy (ART) late in the course of HIV disease by gender



Indicator 14 Proportion of individuals starting antiretroviral therapy (ART) late in the course of HIV disease

Interpretations & Comments	The proportion of individuals initiating therapy late in the disease course has historically seen a modest decline. Over the course of 2011 this decline continued in and rates have stabilized for the most part in the first half of 2012. NI continues to experience large fluctuations associated with the small sample size. Trends have historically been similar for both sexes and the modest divergence seen in the last quarter has disappeared.
Description of Measure	Percentage of individuals starting ART who have cd4 cell counts below 200 cells/mL.
Significance	Current clinical guidelines are complex in terms of eligibility for ART and rely on an algorithm which takes into account cd4, viral load, concomitant illness or other morbidities including laboratory findings, other medications and their safety profiles as well as the lifestyle or personal challenges of the individual. A cd4 cell count of <200 cells/mL however, represents severe HIV disease and a point at which all individuals should be accessing therapy.
Data Source(s)	British Columbia Center for Excellence Drug Treatment Program Database
Calculation Method	Denominator: Total number of distinct individuals who are initiating ART in the time period of interest. Numerator: Total number of individuals from the denominator with cd4 cell counts lower than 200 cells/mL
Limitations	CD4 count is only one measure of treatment eligibility. Therefore, there are likely to be individuals who are eligible for treatment on other grounds but whose cd4 count is greater than 200 and will not be captured in the numerator in this estimate.
Notes	
Revisions	

Indicator 18: Proportion of individuals with a new HIV diagnosis who are tested for syphilis within three months of HIV diagnosis

Target:	>95%	
Actual:	VAN: 73.3% in 2011 Q3 & Q4	NI: 33.3% in 2011 Q3 & Q4

Figure 18.1 Proportion of individuals with a new HIV diagnosis who are tested for syphilis within 3 months of HIV diagnosis by HSDA




Figure 18.2 Proportion of individuals with a new HIV diagnosis who are tested for syphilis within 3 months of HIV diagnosis by HSDA – Males




Figure 18.3 Proportion of individuals with a new HIV diagnosis who are tested for syphilis within 3 months of HIV diagnosis by HSDA – Females





Figure 18.4 Proportion of individuals with a new HIV diagnosis who are tested for syphilis within 3 months of HIV diagnosis – by Aboriginal status



Indicator 18 Proportion of individuals with a new HIV diagnosis who are tested for syphilis within 3 months of HIV diagnosis

Interpretations & Comments	In 2011 Q3&Q4, the proportion of individuals with a new HIV diagnosis who are tested for syphilis within 3 months of diagnosis increased in Vancouver HSDA, was stable in other HSDAs, and decreased in Northern Interior HSDA. Similar trends were observed for males while trends in females were more variable; the magnitude of this proportion is lower for females compared to males. The proportion in Aboriginal persons is variable.
Description of Measure	The percent of individuals with a new diagnosis of HIV who have a syphilis test within 3 months of their HIV diagnosis date.
Significance	Testing for sexually transmitted infections including syphilis is recommended routinely for individuals with HIV upon entry into HIV-related primary care and by public health during follow-up of new positive HIV tests. Measuring the proportion of individuals with a new diagnosis of HIV who have a syphilis test within 3 months after the date of HIV diagnosis may be a proxy for entry into HIV-related primary care and success of public health follow-up.
Data Source(s)	<ul style="list-style-type: none"> Provincial HIV/AIDS surveillance database at BCCDC. Misys Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory (PHSA).
Calculation Method	<ul style="list-style-type: none"> Based on a direct match of identifiers for individuals with a new positive HIV test and individuals undergoing syphilis testing. Denominator: All unique individuals with a new HIV diagnosis. Numerator: Number of unique individuals with a new HIV diagnosis who have a syphilis test within 14 days before or 3 months after the date of HIV diagnosis. Allocation by HSDA is based on address of individual with new HIV diagnosis, or if unknown, address of ordering clinician or clinic. Unit of analysis is the percentage of individuals with a new HIV diagnosis who are tested within 3 months for syphilis, per six months.
Limitations	<ul style="list-style-type: none"> Individuals who test for HIV using different identifiers (e.g., initials, pseudonyms, non-nominally) than are used for syphilis testing will not be included in the numerator. <ul style="list-style-type: none"> POC HIV test data and HIV test data from another laboratory are not included in the data linkage. This indicator has similar limitations as 6b, as the linkage involves two distinct databases and the potential for misclassification is higher. Due to data quality limitations this indicator underestimates the true proportion.
Notes	
Revisions	<ul style="list-style-type: none"> Indicator debuted. (Oct 2010) Improvement to the method for data analysis has revised the values of this indicator slightly from the November 10, 2010 report. (Jan 2011) Breakdown by Aboriginal status included (Oct 2011)

Indicator 21: Percentage of HIV-infected individuals who are tested for genotypic antiretroviral drug resistance prior to starting antiretroviral therapy (ART)

Target:	Increase to >95%	
Actual:	VAN: 88.89%	NI: 100.00%

Figure 21.1 Percentage of HIV-infected individuals who are tested for genotypic antiretroviral drug resistance prior to starting antiretroviral therapy (ART) by HSDA





Figure 21.2 Percentage of HIV-infected individuals who are tested for genotypic antiretroviral drug resistance prior to starting antiretroviral therapy (ART) by gender



Indicator 21 Percentage of HIV-infected individuals who are tested for genotypic antiretroviral drug resistance prior to starting antiretroviral therapy (ART)

Interpretations & Comments	Estimates of the proportion of individuals receiving genotypic testing have remained relatively steady throughout 2011 and the first half of 2012. The overall trend is towards rates near goal levels for both men and women.
Description of Measure	Percentage of HIV positive individuals who receive laboratory testing for genotypic drug resistance before they begin antiretroviral therapy.
Significance	Over time individuals exposed to ART can develop strains of HIV that are resistant to some or all of the drugs in a given therapy regimen. When this happens the efficacy of the drugs declines and the drug regimen must be changed. People with resistant virus can pass along these resistant virus strains so that those they infect actually have drug resistance even though they have never taken antiretroviral drugs. Therefore, it is important to conduct genotype testing on those who have never been exposed to ART but who are initiating therapy. The purpose of this is to establish whether the patient is harboring drug resistant strains of the HI virus so that the therapy can be tailored to suit the patient's needs. Testing typically includes resistance to nucleoside reverse transcriptase inhibitors (NRTI), non-nucleoside reverse transcriptase inhibitors (NNRTI), M18, and protease inhibitor (PI) classes of therapy.
Data Source(s)	British Columbia Center for Excellence Drug Treatment Program Database
Calculation Method	Percentage of all those initiating first therapy who have prior genotype testing. Denominator: All individuals who initiated first ever antiretroviral therapy Numerator: All those in the denominator that have had at least one resistance profile conducted prior to therapy start date.
Limitations	Viral load must be ≥ 250 copies/mL for testing to be conducted. Prior to January 1, 2002 pVL needed to be $\geq 1,000$ copies/mL.
Notes	
Revisions	

Indicator 22: Percentage of individuals starting antiretroviral therapy (ART) who achieve HIV plasma viral load (pVL) of <200 copies/mL within nine months of therapy initiation

Target: Increase to >95%

Actual: VAN: 91.55%

NI: 75.00%

Figure 22.1 Percentage of individuals starting ART who achieve HIV plasma viral load (pVL) of < 200 copies/mL within 9 months of therapy initiation by HSDA





Figure 22.2 Percentage of individuals starting ART who achieve HIV plasma viral load (pVL) of < 200 copies/mL within nine months of therapy initiation by gender



Indicator 22 Percentage of individuals starting antiretroviral therapy (ART) who achieve HIV plasma viral load (pVL) of <200 copies/mL within nine months of therapy initiation

Interpretations & Comments	Rates in Vancouver and Northern Interior continue to fluctuate and fall short of the 95% goal. The past quarter reveals a decline among women, and although rates for women typically have experienced greater variation, a 20% decline over 3 months will require careful monitoring.
Description of Measure	Percentage of individuals initiating first antiretroviral therapy who have a pVL below the limit of detection within the first nine months of ART.
Significance	Plasma viral load is a measure of viral activity assessed by quantifying the amount of virus present in the patient's blood. Lower pVL is associated with reduced disease activity with counts below the limit of detection indicating excellent virus suppression—the ultimate goal of ART. As long as viral suppression is maintained disease progression is curtailed. Individuals receiving appropriate therapy in accordance with clinical guidelines are generally expected to successfully suppress virus within the first six months of treatment. However, imperfect adherence to therapy or resistance due to primary infection with a drug resistant strain of HIV can negatively impact therapy success.
Data Source(s)	British Columbia Center for Excellence Drug Treatment Program Database
Calculation Method	Denominator: All individuals initiating first ever ART. Numerator: Of individuals in the denominator, those who had two consecutive pVL measures <200 copies/mL both taken after therapy start and at least one of which is taken within the first nine months of treatment
Limitations	Can be confounded by patient-related factors including adherence. Prior to February 1 st , 2008 the lowest limit of detection was considered to be pVL<50copies/mL. Since that time a new laboratory technique has been adopted to quantify pVL. This method is less accurate at low pVL levels and currently a pVL<200 is considered to represent complete suppression.
Notes	
Revisions	For the first three quarters of 2010 reports were presented using pVL<50 copies/mL to be consistent with older testing techniques. From the last quarter of 2010 forward reporting will use the new standard of 200 copies/mL. Starting in the second quarter of 2011 the indicator was revised from suppression within the first six months to the first nine months after therapy initiation. This change was enacted because a review of current data found that the use of the six-month timeframe yielded an underestimate of treatment effectiveness due to the automatic inclusion as “failures” of those who did not receive a second test in the six month window.

Indicator 23*: Percentage of individuals who initiated antiretroviral therapy (ART) with a recommended therapy regimen (among those with no drug resistance)

Target: Increase to >95%

Actual: VAN: 72.22%

NI: 100.00%

Figure 23.1 Percentage of individuals who initiated antiretroviral therapy (ART) with a recommended therapy regimen (among those with no drug resistance) by HSDA





Figure 23.2 Percentage of individuals who initiated antiretroviral therapy (ART) with a recommended therapy regimen (among those with no drug resistance) by gender



Indicator 23 Percentage of individuals who initiated antiretroviral therapy (ART) with a recommended therapy regimen (among those with no drug resistance)

Interpretations & Comments	In 2012 thus far rates continue to fluctuate around the 75% mark, a distant 20% from the goal level. Both sexes continue to follow similar patterns although women are, once again, experiencing a drift downwards in comparison to men.
Description of Measure	Percentage of individuals who are starting first ever ART and who have been shown to have no drug resistance who initiate therapy with one of the therapy regimens recommended for those who have never been on therapy and who do not have any drug resistance.
Significance	As described in Indicator 21, resistance testing is an important precursor to treatment. Drug resistance complicates treatment and limits treatment options. Individuals without drug resistance have the option of using, and should be prescribed, the most simple and effective therapy options. Currently 8 options are recommended for people who are new to treatment and who do not have drug resistance.
Data Source(s)	British Columbia Center for Excellence Drug Treatment Program Database
Calculation Method	<i>Denominator:</i> All individuals initiating first ever ART who had drug resistance testing prior to ART start date, which documented no resistance to any of nucleoside reverse transcriptase inhibitors (NRTI), non-nucleoside reverse transcriptase inhibitors (NNRTI), M18, and protease inhibitor (PI) classes of therapy. <i>Numerator:</i> Individuals in the denominator who initiated first ever therapy with one of the eight therapy regimens recommended.
Limitations	Patients may have specific contraindications other than resistance and these data are not completely captured.
Notes*	THIS INDICATOR IS UNDER REVIEW TO REFLECT CHANGES IN CURRENTLY RECOMMENDED THERAPY OPTIONS. Q2 FIGURES MAY BE UPDATED IN Q3 TO REFLECT THESE CHANGES AND ARE PROVIDED HERE IN THE INTEREST OF CONSISTENCY ONLY.
Revisions	From study start through Q2 of 2012, the options for first line therapy were: <ul style="list-style-type: none"> • Lamivudine/lopinavir+ritonavir/tenofovir • Lamivudine/efavirenz/tenofovir • Lamivudine/nevirapine/tenofovir • Lamivudine/tenofovir/ritonavir boosted atazanavir • Lopinavir+ritonavir/tenofovir/emtricitabine • Efavirenz/tenofovir/emtricitabine • Nevirapine/tenofovir/emtricitabine • Tenofovir/ritonavir boosted atazanavir/emtricitabine

Indicator 24: Percentage of individuals on antiretroviral therapy (ART) that achieve annual prescription refill adherence of >95%

Target:	Increase	
Actual:	VAN: 77.41%	NI: 59.26%

Figure 24.1 Percentage of individuals on antiretroviral therapy (ART) that achieve annual prescription refill adherence of >95% by HSDA





Figure 24.2 Percentage of individuals on antiretroviral therapy (ART) that achieve annual prescription refill adherence of >95% by gender




Indicator 24 Percentage of individuals on antiretroviral therapy (ART) that achieve annual prescription refill adherence of >95%

Interpretations & Comments	The trend for all HSDA remains stable and, while the NI experiences lower rates of adherence overall and broader variation in estimates due to low numbers, rates in this HSDA also remain relatively consistent and have recently experienced an upward trend. Women continue to have lower rates of high adherence although this gap appears to be closing slightly.
Description of Measure	Percentage of individuals starting ART that pick up at least 95% of their prescribed medication over the first year of therapy.
Significance	For therapy to be effective the prescribed drugs must be taken as directed. One of the primary reasons for treatment failure is incomplete adherence (missed drug doses). In fact, levels of adherence of around 95% have been correlated with sustained virologic suppression, fewer hospitalizations, and reduced rates of drug resistance.
Data Source(s)	British Columbia Center for Excellence Drug Treatment Program Database
Calculation Method	<i>Denominator:</i> All individuals prescribed ART <i>Numerator:</i> All individuals in the denominator who have at least 95% adherence over the past full year of therapy Adherence is calculated as: <i>Denominator:</i> 365 days <i>Numerator:</i> Total number of days covered by prescriptions filled (i.e., picked up by the patient) from start date of ART to day 365.
Limitations	This measure is a proxy for adherence to ART. Adherence will be overestimated if prescriptions are filled but medication is not taken. Missed medication pick-ups may be a result of medically ordered temporary treatment interruptions rather than patient non-adherence. Patients may have stockpiles of medication at home from prior years and so may miss pickups yet remain adherent.
Notes	
Revisions	

Indicator 25: Number of physicians initiating therapy or providing HIV-related care to patients on antiretroviral therapy (ART)

Target:	Increase	
Actual:	VAN: 201	NI: 25

Figure 25.1 Number of physicians initiating antiretroviral therapy (ART) by HSDA



Indicator 25 Number of physicians initiating antiretroviral therapy (ART)

Interpretations & Comments	The total number of prescribing physicians remains relatively stable in the long term with modest but consistent gains over time in most HSDA. The situation in the Northern HSDA with few physicians providing HIV therapy-related care remains precarious.
Description of Measure	The number of doctors who are initiating HIV-related ART.
Significance	Access to high quality care close to home is of great concern for patients. The total number of physicians in a given geographic area successfully prescribing ART as either the enrolling or follow-up physician is one important measure of access for patients to HIV care.
Data Source(s)	British Columbia Center for Excellence Drug Treatment Program Database
Calculation Method	Simple count of the total number of physicians in the geographic area of interest who are successfully initiating ART among HIV positive individuals. Successful prescription is defined as having at least one prescription for ART filled for a patient not previously on ART. The unit of analysis is the unique physician.
Limitations	
Notes	
Revisions	

Indicator 26: Percentage of individuals on antiretroviral therapy (ART) who experience a serious adverse drug reaction (ADR)

Target:	Maintain <0.5%	
Actual:	VAN: 0.09%	NI: 0.00%

Figure 26.1 Percentage of individuals on antiretroviral therapy (ART) who experience a serious adverse drug reaction (ADR) by HSDA





Figure 26.2 Percentage of individuals on antiretroviral therapy (ART) who experience a serious adverse drug reaction (ADR) by gender



Indicator 26 Percentage of individuals on antiretroviral therapy (ART) who experience a serious adverse drug reaction (ADR)

Interpretations & Comments	<p>The trend remains towards very low ADR rates. While women may experience (or their physicians report) greater rates of ADR this gap has closed over the past three quarters.</p> <p>Due to the small number of events trends in this indicator must be interpreted with caution- particularly in the Northern HSDA where a single case can cause a dramatic spike in comparative quarterly rates.</p>
Description of Measure	Percentage of individuals on ART who have a serious negative reaction to an ART drug.
Significance	<p>Most medications can be associated with adverse reactions. Serious adverse drug events in HIV therapy cover a wide range of problems in various organ systems and are defined as reactions that are potentially life-threatening or which lead to hospitalization or death. Monitoring for ADRs in the general population of ART users is important because the clinical trials in which drug testing is conducted usually include relatively few patients followed over a comparatively short time period. Therefore, trials may not identify ADR if they are very rare or are a result of very long exposure. Fortunately, the risk of a serious ADR in response to antiretroviral drugs is very low.</p>
Data Source(s)	British Columbia Center for Excellence Drug Treatment Program Database
Calculation Method	<p><i>Denominator:</i> Total number of distinct individuals who are taking ART and any given time in the time period of interest.</p> <p><i>Numerator:</i> Number of serious adverse events over the time period of interest.</p>
Limitations	Reporting of adverse drug reactions, even serious ones, is voluntary- relying on physician report. Moreover, those that are reported are not confirmed or substantiated independently and it remains unknown whether factors other than ART drugs may be responsible or partially responsible for the adverse event.
Notes	
Revisions	

Indicator 28: Incidence of resistance to any retroviral drug

Target:	Decrease	
Actual:	VAN: 0.05%	NI: 0.31%

Figure 28.1 Incidence of resistance to any antiretroviral drug by HSDA





Figure 28.2 Incidence of resistance to any antiretroviral drug by gender



Indicator 28 Incidence of resistance to any antiretroviral drug

Interpretations & Comments	All HSDA have consistently low rates of incident drug resistance throughout 2010/2011 and the first half of 2012. The slightly higher rates among women observed historically have resolved to a large degree in the past six months.
Description of Measure	Counts new cases of antiretroviral drug resistance occurring over the time period of interest among all individuals taking antiretroviral therapy.
Significance	One goal of the STOP HIV pilot is to reduce transmission of drug-resistant HIV strains. The lower the incidence of resistance and the fewer people with HIV harboring resistant viral strains, the more successful these efforts will be.
Data Source(s)	British Columbia Center for Excellence Drug Treatment Program Database and genotypic testing database held at the British Columbia Center for Excellence laboratory
Calculation Method	<i>Numerator:</i> Number of new (excludes previously identified resistance) cases of drug resistance detected in each quarter <i>Denominator:</i> Total number of person-months of antiretroviral exposure in the quarter.
Limitations	This indicator show trends in the detection of resistance, however temporal trends in the frequency of resistance testing (increasing rates over time) may confound trends in the actual occurrence of resistance. Genotyping can only be conducted for individuals with pVL >= 250 copies/mL (although this may be of little clinical relevance).
Notes	
Revisions	

Indicator 29: Proportion of individuals on antiretroviral therapy who change antiretroviral drug treatment

Target: Decrease

Actual: VAN: 3.70%

NI: 2.46%

Figure 29.1 Proportion of individuals on antiretroviral therapy who change antiretroviral drug treatment by HSDA





Figure 29.2 Proportion of individuals on antiretroviral therapy who change antiretroviral drug treatment by gender



Indicator 29 Proportion of individuals on antiretroviral therapy who change antiretroviral drug treatment

Interpretations & Comments	The trend remains steady with consistently low rates and over the past three quarters in all HSDA and among both sexes.
Description of Measure	The percentage of all individuals on antiretroviral therapy who change their therapeutic regimen over the course of the time period of interest.
Significance	Changes in therapy regimen occur most commonly as a result of drug intolerance, adverse drug reactions or treatment failure. By counting the occurrence of regimen change and identifying the reasons for these changes a broader and more inclusive estimate of the safety of antiretroviral therapies can be made. Please see limitations of adverse drug event reporting under Indicator 26.
Data Source(s)	British Columbia Center for Excellence Drug Treatment Program Database
Calculation Method	<i>Numerator:</i> Total number of regimen changes, where a regimen change is defined as a class change in the NNRTI or PI component of the therapy regimen. <i>Denominator:</i> Total number of individuals on antiretroviral therapy.
Limitations	The reason for change is often not well recorded and the indicator relies heavily on exclusion of treatment failure as the reason for therapy change.
Notes	
Revisions	

Data Tables

Table 1.1 Number of HIV test episodes by HSDA

Quarter	Vancouver HSDA	Northern Interior HSDA	Other HSDAs	All HSDAs
2006 Q2	12,547	1,314	26,399	40,260
2006 Q3	12,778	1,415	27,531	41,724
2006 Q4	12,337	1,271	26,581	40,189
2007 Q1	13,673	1,572	30,584	45,829
2007 Q2	12,475	1,389	27,215	41,079
2007 Q3	13,148	1,365	27,276	41,789
2007 Q4	12,844	1,333	27,593	41,770
2008 Q1	14,127	1,446	29,448	45,021
2008 Q2	14,232	1,417	29,356	45,005
2008 Q3	14,471	1,428	28,899	44,798
2008 Q4	13,743	1,363	28,869	43,975
2009 Q1	15,231	1,518	30,551	47,300
2009 Q2	14,418	1,331	27,961	43,710
2009 Q3	14,833	1,343	27,967	44,143
2009 Q4	13,844	1,247	26,486	41,577
2010 Q1	15,262	1,459	29,670	46,391
2010 Q2	14,797	1,278	28,086	44,161
2010 Q3	15,098	1,341	28,249	44,688
2010 Q4	15,727	1,310	28,494	45,531
2011 Q1	17,023	1,525	30,210	48,758
2011 Q2	16,603	1,365	27,620	45,588
2011 Q3	22,435	1,433	29,155	53,023
2011 Q4	20,204	1,523	29,208	50,935
2012 Q1	22,837	1,795	32,109	56,741
2012 Q2	22,512	1,583	30,824	54,919

Table 1.2 Number of HIV test episodes by HSDA – Males

Quarter	Vancouver HSDA	Northern Interior HSDA	Other HSDAs	All HSDAs
2006 Q2	5,602	435	9,101	15,138
2006 Q3	5,585	456	9,479	15,520
2006 Q4	5,426	429	9,135	14,990
2007 Q1	6,161	502	10,558	17,221
2007 Q2	5,544	446	9,338	15,328
2007 Q3	5,929	429	9,198	15,556
2007 Q4	5,712	385	9,391	15,488
2008 Q1	6,323	503	10,068	16,894
2008 Q2	6,406	451	10,302	17,159
2008 Q3	6,562	512	9,951	17,025
2008 Q4	6,168	452	10,167	16,787
2009 Q1	7,077	525	10,505	18,107
2009 Q2	6,547	439	9,556	16,542
2009 Q3	6,798	454	9,491	16,743
2009 Q4	6,307	356	8,798	15,461
2010 Q1	7,034	535	10,075	17,644
2010 Q2	6,897	435	9,809	17,141
2010 Q3	6,932	436	9,627	16,995
2010 Q4	6,901	398	9,733	17,032
2011 Q1	7,398	472	10,408	18,278
2011 Q2	6,759	469	9,428	16,656
2011 Q3	7,776	468	10,000	18,244
2011 Q4	8,205	475	10,151	18,831
2012 Q1	9,756	599	11,325	21,680
2012 Q2	9,833	567	10,754	21,154

Table 1.3 Number of HIV test episodes by HSDA – Females

Quarter	Vancouver HSDA	Northern Interior HSDA	Other HSDAs	All HSDAs
2006 Q2	6,728	855	17,005	24,588
2006 Q3	6,950	897	17,581	25,428
2006 Q4	6,630	806	16,972	24,408
2007 Q1	7,222	1,008	19,361	27,591
2007 Q2	6,665	882	17,349	24,896
2007 Q3	6,983	899	17,632	25,514
2007 Q4	6,932	911	17,809	25,652
2008 Q1	7,565	919	19,069	27,553
2008 Q2	7,553	915	18,702	27,170
2008 Q3	7,618	891	18,639	27,148
2008 Q4	7,315	889	18,445	26,649
2009 Q1	7,806	957	19,729	28,492
2009 Q2	7,416	869	18,160	26,445
2009 Q3	7,578	869	18,201	26,648
2009 Q4	7,160	877	17,481	25,518
2010 Q1	7,791	908	19,305	28,004
2010 Q2	7,487	832	18,003	26,322
2010 Q3	7,754	890	18,448	27,092
2010 Q4	7,536	851	18,560	26,947
2011 Q1	8,132	1,018	19,587	28,737
2011 Q2	7,454	835	17,973	26,262
2011 Q3	8,309	901	18,834	28,044
2011 Q4	9,206	976	18,857	29,039
2012 Q1	10,806	1,092	20,562	32,460
2012 Q2	10,895	927	19,799	31,621

Table 1.4 Number of POC HIV tests by HSDA

	Vancouver HSDA	Northern Interior HSDA
2006 Q2		
2006 Q3		
2006 Q4		
2007 Q1		
2007 Q2		
2007 Q3		
2007 Q4		
2008 Q1		
2008 Q2		
2008 Q3		
2008 Q4		
2009 Q1		
2009 Q2		
2009 Q3		
2009 Q4		
2010 Q1	312	0
2010 Q2	396	0
2010 Q3	503	0
2010 Q4	915	45
2011 Q1	1059	19
2011 Q2	1971	49
2011 Q3	5912	54
2011 Q4	2379	52
2012 Q1	1987	83
2012 Q2	1552	74

Table 2.1 Population HIV testing rate by HSDA

Year	Vancouver HSDA		Northern Interior HSDA		Other HSDAs		All HSDAs	
	HIV Test	Rate	HIV Test	Rate	HIV Test	Rate	HIV Test	Rate
2006	38,940	6,356.2	5,227	3,720.9	109,850	3,147.1	154,017	3,629.4
2007	37,401	5,998.2	5,284	3,748.4	114,728	3,236.2	157,413	3,652.6
2008	40,946	6,500.4	5,209	3,667.2	118,085	3,269.3	164,240	3,746.5
2009	42,634	6,627.8	5,028	3,521.9	114,858	3,126.0	162,520	3,643.7
2010	43,010	6,524.4	4,944	3,438.4	116,876	3,135.1	164,830	3,637.9
2011	46,305	6,965.0	5,248	3,641.4	119,075	3,154.2	170,628	3,722.2

Table 2.2 Population HIV testing rate by HSDA – Males

Year	Vancouver HSDA		Northern Interior HSDA		Other HSDAs		All HSDAs	
	HIV Test	Rate	HIV Test	Rate	HIV Test	Rate	HIV Test	Rate
2006	17,483	5,760.0	1,716	2,399.0	37,946	2,197.1	57,145	2,718.4
2007	16,940	5,479.6	1,680	2,339.8	39,550	2,254.0	58,170	2,723.8
2008	18,440	5,899.7	1,784	2,465.0	41,506	2,320.8	61,730	2,840.3
2009	19,415	6,078.1	1,635	2,247.2	39,668	2,179.7	60,718	2,744.8
2010	19,828	6,056.4	1,673	2,283.6	40,807	2,210.4	62,308	2,773.2
2011	21,199	6,419.6	1,775	2,419.2	42,051	2,250.0	65,025	2,861.3

Table 2.3 Population HIV testing rate by HSDA – Females

Year	Vancouver HSDA		Northern Interior HSDA		Other HSDAs		All HSDAs	
	HIV Test	Rate	HIV Test	Rate	HIV Test	Rate	HIV Test	Rate
2006	20,524	6,761.9	3,345	4,676.4	70,210	4,065.2	94,079	4,475.4
2007	19,520	6,314.1	3,419	4,761.7	73,224	4,173.1	96,163	4,502.8
2008	21,469	6,868.8	3,307	4,569.3	75,372	4,214.3	100,148	4,607.9
2009	21,616	6,767.2	3,303	4,539.8	74,187	4,076.4	99,106	4,480.2
2010	21,600	6,597.7	3,216	4,389.8	75,160	4,071.3	99,976	4,449.8
2011	23,441	7,098.5	3,418	4,658.5	76,371	4,086.3	103,230	4,542.5

Table 3.1 Number of new HIV diagnoses by HSDA – Allocated by RESIDENCE

Quarter	Vancouver HSDA	Northern Interior HSDA	Other HSDAs	All HSDAs
2006 Q1	51	6	39	96
2006 Q2	50	3	40	93
2006 Q3	36	5	47	88
2006 Q4	42	5	37	84
2007 Q1	66	5	39	110
2007 Q2	51	6	45	102
2007 Q3	35	9	43	87
2007 Q4	40	3	49	92
2008 Q1	54	0	43	97
2008 Q2	40	4	36	80
2008 Q3	40	3	45	89
2008 Q4	40	3	36	79
2009 Q1	45	4	53	102
2009 Q2	34	4	46	84
2009 Q3	38	2	38	78
2009 Q4	34	6	33	73
2010 Q1	36	1	36	73
2010 Q2	40	0	41	81
2010 Q3	34	5	39	78
2010 Q4	38	2	28	68
2011 Q1	29	0	24	53
2011 Q2	46	5	31	82
2011 Q3	51	4	33	88
2011 Q4	36	4	25	65
2012 Q1	43	3	25	72
2012 Q2	32	3	22	57

Table 3.2 Number of new HIV diagnoses by HSDA – Allocated by ORDERING CLINICIAN

Quarter	Vancouver HSDA	Northern Interior HSDA	Other HSDAs	All HSDAs
2006 Q1	57	6	33	96
2006 Q2	52	3	38	93
2006 Q3	47	5	36	88
2006 Q4	45	4	35	84
2007 Q1	67	5	38	110
2007 Q2	57	5	40	102
2007 Q3	41	10	36	87
2007 Q4	51	4	37	92
2008 Q1	62	0	35	97
2008 Q2	49	4	27	80
2008 Q3	44	4	41	89
2008 Q4	47	4	28	79
2009 Q1	52	4	46	102
2009 Q2	44	4	36	84
2009 Q3	41	3	34	78
2009 Q4	39	7	27	73
2010 Q1	42	1	30	73
2010 Q2	46	0	35	81
2010 Q3	40	5	33	78
2010 Q4	41	2	25	68
2011 Q1	32	0	21	53
2011 Q2	43	5	34	82
2011 Q3	62	4	22	88
2011 Q4	40	4	21	65
2012 Q1	46	3	23	72
2012 Q2	39	4	14	57

Table 3.3 Number of new HIV diagnoses by gender, BC

Quarter	Male	Female	Other	All
2006 Q1	81	15	0	96
2006 Q2	74	19	0	93
2006 Q3	67	21	0	88
2006 Q4	67	17	0	84
2007 Q1	88	21	1	110
2007 Q2	79	21	2	102
2007 Q3	65	22	0	87
2007 Q4	71	21	0	92
2008 Q1	83	14	0	97
2008 Q2	64	16	0	80
2008 Q3	75	14	0	89
2008 Q4	64	15	0	79
2009 Q1	81	21	0	102
2009 Q2	68	16	0	84
2009 Q3	61	17	0	78
2009 Q4	56	17	0	73
2010 Q1	56	17	0	73
2010 Q2	61	20	0	81
2010 Q3	64	14	0	78
2010 Q4	57	11	0	68
2011 Q1	46	6	1	53
2011 Q2	63	19	0	82
2011 Q3	77	11	0	88
2011 Q4	58	7	0	65
2012 Q1	62	10	0	72
2012 Q2	47	10	0	57

Other = Transgender + Gender Unknown

Table 3.4 Number of new HIV diagnoses by Aboriginal status, BC

Quarter	Aboriginal	Non Aboriginal	Unknown Ethnicity	All
2006 Q1	12	80	4	96
2006 Q2	17	74	2	93
2006 Q3	15	68	5	88
2006 Q4	12	66	6	84
2007 Q1	20	84	6	110
2007 Q2	16	78	8	102
2007 Q3	13	73	1	87
2007 Q4	14	72	6	92
2008 Q1	12	82	3	97
2008 Q2	10	66	4	80
2008 Q3	7	76	6	89
2008 Q4	17	56	6	79
2009 Q1	17	81	4	102
2009 Q2	13	67	4	84
2009 Q3	13	61	4	78
2009 Q4	14	55	4	73
2010 Q1	7	65	1	73
2010 Q2	13	64	4	81
2010 Q3	13	63	2	78
2010 Q4	6	57	5	68
2011 Q1	8	44	1	53
2011 Q2	15	63	4	82
2011 Q3	10	71	7	88
2011 Q4	12	51	2	65

Unknown: ethnicity not stated

Table 4.1 Rate of new AIDS case reports by HSDA

Year	Vancouver HSDA		Northern Interior HSDA		Other HSDAs		All HSDAs	
	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
2006	42	6.9	3	2.1	65	1.9	110	2.6
2007	54	8.7	4	2.8	49	1.4	107	2.5
2008	54	8.6	1	0.7	55	1.5	110	2.5
2009	35	5.4	5	3.5	45	1.2	85	1.9
2010	20	3.1	0	0	36	1.0	56	1.2

Table 4.2 Rate of new AIDS case reports, by gender, BC

Year	Male		Female		Other		All	
	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
2006	94	4.5	16	0.7	0	---	110	2.6
2007	94	4.4	13	0.6	0	---	107	2.5
2008	87	4	23	1	0	---	110	2.5
2009	70	3.2	15	0.7	0	---	85	1.9
2010	41	1.8	15	0.7	0	---	56	1.2

Other = Transgender + Gender Unknown

Table 5.1 Percentage positivity among persons tested for HIV by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDA			All HSDA		
	First HIV+	HIV Test	%	First HIV+	HIV Test	%	First HIV+	HIV Test	%	First HIV+	HIV Test	%
2006 Q2	62	12,431	0.50%	3	1,302	0.23%	43	26,180	0.16%	108	39,913	0.27%
2006 Q3	59	12,667	0.47%	5	1,405	0.36%	43	27,288	0.16%	107	41,360	0.26%
2006 Q4	64	12,225	0.52%	5	1,258	0.40%	44	26,383	0.17%	113	39,866	0.28%
2007 Q1	80	13,540	0.59%	5	1,561	0.32%	49	30,326	0.16%	134	45,427	0.29%
2007 Q2	68	12,342	0.55%	7	1,378	0.51%	49	26,997	0.18%	124	40,717	0.30%
2007 Q3	57	13,001	0.44%	10	1,352	0.74%	45	27,063	0.17%	112	41,416	0.27%
2007 Q4	62	12,727	0.49%	5	1,315	0.38%	52	27,392	0.19%	119	41,434	0.29%
2008 Q1	79	13,964	0.57%	1	1,430	0.07%	49	29,179	0.17%	129	44,573	0.29%
2008 Q2	68	14,079	0.48%	6	1,408	0.43%	30	29,094	0.10%	104	44,581	0.23%
2008 Q3	55	14,298	0.38%	3	1,412	0.21%	52	28,629	0.18%	110	44,339	0.25%
2008 Q4	59	13,615	0.43%	4	1,344	0.30%	41	28,648	0.14%	104	43,607	0.24%
2009 Q1	65	15,053	0.43%	4	1,503	0.27%	58	30,291	0.19%	127	46,847	0.27%
2009 Q2	60	14,257	0.42%	4	1,322	0.30%	49	27,712	0.18%	113	43,291	0.26%
2009 Q3	52	14,674	0.35%	4	1,326	0.30%	41	27,725	0.15%	97	43,725	0.22%
2009 Q4	49	13,714	0.36%	7	1,240	0.56%	40	26,274	0.15%	96	41,228	0.23%
2010 Q1	61	15,062	0.40%	1	1,440	0.07%	34	29,385	0.12%	96	45,887	0.21%
2010 Q2	53	14,624	0.36%	0	1,268	0.00%	42	27,870	0.15%	95	43,762	0.22%
2010 Q3	54	14,900	0.36%	4	1,330	0.30%	45	28,012	0.16%	103	44,242	0.23%
2010 Q4	51	14,638	0.35%	2	1,250	0.16%	28	28,239	0.10%	81	44,127	0.18%
2011 Q1	40	15,731	0.25%	0	1,491	0.00%	26	29,929	0.09%	66	47,151	0.14%
2011 Q2	61	14,454	0.42%	4	1,306	0.31%	43	27,302	0.16%	108	43,062	0.25%
2011 Q3	88	16,285	0.54%	5	1,357	0.37%	37	28,752	0.13%	130	46,394	0.28%
2011 Q4	58	17,605	0.33%	4	1,450	0.28%	30	28,912	0.10%	92	47,967	0.19%
2012 Q1	62	20,568	0.30%	8	1,691	0.47%	34	31,703	0.11%	104	53,962	0.19%
2012 Q2	54	20,697	0.26%	6	1,483	0.40%	21	30,393	0.07%	81	52,573	0.15%

Table 5.2 Percentage positivity among persons tested for HIV by gender, BC

Quarter	Male			Female			Unknown			All		
	First HIV+	HIV Test	%	First HIV+	HIV Test	%	First HIV+	HIV Test	%	First HIV+	HIV Test	%
2006 Q2	77	15,001	0.51%	27	24,381	0.11%	4	531	0.75%	108	39,913	0.27%
2006 Q3	78	15,391	0.51%	24	25,195	0.10%	5	774	0.65%	107	41,360	0.26%
2006 Q4	90	14,859	0.61%	22	24,220	0.09%	1	787	0.13%	113	39,866	0.28%
2007 Q1	104	17,046	0.61%	28	27,374	0.10%	2	1,007	0.20%	134	45,427	0.29%
2007 Q2	99	15,188	0.65%	23	24,682	0.09%	2	847	0.24%	124	40,717	0.30%
2007 Q3	84	15,405	0.55%	27	25,294	0.11%	1	717	0.14%	112	41,416	0.27%
2007 Q4	91	15,366	0.59%	25	25,436	0.10%	3	632	0.47%	119	41,434	0.29%
2008 Q1	109	16,705	0.65%	19	27,291	0.07%	1	577	0.17%	129	44,573	0.29%
2008 Q2	81	16,990	0.48%	23	26,918	0.09%	0	673	0.00%	104	44,581	0.23%
2008 Q3	87	16,843	0.52%	21	26,870	0.08%	2	626	0.32%	110	44,339	0.25%
2008 Q4	81	16,659	0.49%	22	26,412	0.08%	1	536	0.19%	104	43,607	0.24%
2009 Q1	98	17,917	0.55%	27	28,234	0.10%	2	696	0.29%	127	46,847	0.27%
2009 Q2	93	16,370	0.57%	20	26,201	0.08%	0	720	0.00%	113	43,291	0.26%
2009 Q3	75	16,554	0.45%	21	26,418	0.08%	1	753	0.13%	97	43,725	0.22%
2009 Q4	73	15,322	0.48%	22	25,309	0.09%	1	597	0.17%	96	41,228	0.23%
2010 Q1	75	17,429	0.43%	21	27,719	0.08%	0	739	0.00%	96	45,887	0.21%
2010 Q2	73	16,958	0.43%	22	26,109	0.08%	0	695	0.00%	95	43,762	0.22%
2010 Q3	81	16,829	0.48%	22	26,815	0.08%	0	598	0.00%	103	44,242	0.23%
2010 Q4	70	16,853	0.42%	11	26,706	0.04%	0	568	0.00%	81	44,127	0.18%
2011 Q1	59	18,077	0.33%	6	28,447	0.02%	1	627	0.16%	66	47,151	0.14%
2011 Q2	84	16,461	0.51%	24	26,017	0.09%	0	584	0.00%	108	43,062	0.25%
2011 Q3	108	18,028	0.60%	22	27,732	0.08%	0	634	0.00%	130	46,394	0.28%
2011 Q4	76	18,624	0.41%	15	28,768	0.05%	1	575	0.17%	92	47,967	0.19%
2012 Q1	84	21,413	0.39%	20	32,124	0.06%	0	425	0.00%	104	53,962	0.19%
2012 Q2	68	20,897	0.33%	13	31,297	0.04%	0	379	0.00%	81	52,573	0.15%

Other = Transgender + Gender Unknown

Table 6a.1 Proportion of individuals tested for syphilis who are tested for HIV at the same clinical encounter by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	Syphilis & HIV Test	Syphilis Test	%	Syphilis & HIV test	Syphilis Test	%	Syphilis & HIV test	Syphilis Test	%	Syphilis & HIV test	Syphilis Test	%
2006 Q2	8,587	11,287	76.1%	695	920	75.5%	15,473	19,234	80.4%	24,755	31,441	78.7%
2006 Q3	9,133	11,758	77.7%	747	1,017	73.5%	16,701	20,447	81.7%	26,581	33,222	80.0%
2006 Q4	8,775	11,206	78.3%	718	915	78.5%	16,743	20,208	82.9%	26,236	32,329	81.2%
2007 Q1	9,941	12,511	79.5%	913	1,122	81.4%	19,587	23,225	84.3%	30,441	36,858	82.6%
2007 Q2	9,130	11,430	79.9%	839	1,051	79.8%	17,552	20,961	83.7%	27,521	33,442	82.3%
2007 Q3	9,580	11,782	81.3%	847	1,009	83.9%	18,060	21,491	84.0%	28,487	34,282	83.1%
2007 Q4	9,563	11,872	80.6%	855	1,041	82.1%	18,492	21,772	84.9%	28,910	34,685	83.4%
2008 Q1	10,616	13,215	80.3%	885	1,073	82.5%	20,244	23,656	85.6%	31,745	37,944	83.7%
2008 Q2	10,432	13,268	78.6%	912	1,117	81.6%	19,745	23,211	85.1%	31,089	37,596	82.7%
2008 Q3	10,418	13,141	79.3%	987	1,145	86.2%	19,733	23,083	85.5%	31,138	37,369	83.3%
2008 Q4	10,091	12,474	80.9%	891	1,066	83.6%	19,667	22,891	85.9%	30,649	36,431	84.1%
2009 Q1	11,313	13,761	82.2%	1,001	1,198	83.6%	21,338	25,176	84.8%	33,652	40,135	83.8%
2009 Q2	10,657	13,013	81.9%	899	1,081	83.2%	19,456	23,656	82.2%	31,012	37,750	82.2%
2009 Q3	10,845	13,254	81.8%	886	1,067	83.0%	19,842	24,016	82.6%	31,573	38,337	82.4%
2009 Q4	10,260	12,504	82.1%	865	1,010	85.6%	18,614	22,418	83.0%	29,739	35,932	82.8%
2010 Q1	11,509	13,840	83.2%	938	1,107	84.7%	21,097	25,240	83.6%	33,544	40,187	83.5%
2010 Q2	11,203	13,464	83.2%	837	1,008	83.0%	19,884	23,887	83.2%	31,924	38,359	83.2%
2010 Q3	11,470	13,602	84.3%	948	1,110	85.4%	20,447	24,440	83.7%	32,865	39,152	83.9%
2010 Q4	11,106	13,361	83.1%	845	1,028	82.2%	20,471	24,297	84.3%	32,422	38,686	83.8%
2011 Q1	12,099	14,758	82.0%	1,049	1,224	85.7%	21,718	25,847	84.0%	34,866	41,829	83.4%
2011 Q2	10,904	14,112	77.3%	922	1,074	85.8%	19,617	23,813	82.4%	31,443	38,999	80.6%
2011 Q3	12,279	15,477	79.3%	977	1,149	85.0%	20,921	25,295	82.7%	34,177	41,921	81.5%
2011 Q4	12,447	15,448	80.6%	1,049	1,238	84.7%	20,731	24,817	83.5%	34,227	41,503	82.5%
2012 Q1	14,442	17,807	81.1%	1,186	1,394	85.1%	22,676	27,066	83.8%	38,304	46,267	82.8%
2012 Q2	13,960	17,369	80.4%	1,100	1,287	85.5%	21,260	25,425	83.6%	36,320	44,081	82.4%

Table 6a.2 Proportion of individuals tested for syphilis who are tested for HIV at the same clinical encounter by HSDA – Males

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	Syphilis & HIV Test	Syphilis Test	%	Syphilis & HIV test	Syphilis Test	%	Syphilis & HIV test	Syphilis Test	%	Syphilis & HIV test	Syphilis Test	%
2006 Q2	3,641	5,166	70.5%	129	232	55.6%	3,992	5,535	72.1%	7,762	10,933	71.0%
2006 Q3	3,765	5,293	71.1%	158	279	56.6%	4,396	5,882	74.7%	8,319	11,454	72.6%
2006 Q4	3,681	5,141	71.6%	163	255	63.9%	4,433	5,862	75.6%	8,277	11,258	73.5%
2007 Q1	4,249	5,886	72.2%	178	278	64.0%	5,223	6,668	78.3%	9,650	12,832	75.2%
2007 Q2	3,859	5,257	73.4%	197	301	65.4%	4,765	6,196	76.9%	8,821	11,754	75.0%
2007 Q3	4,117	5,556	74.1%	196	265	74.0%	4,980	6,435	77.4%	9,293	12,256	75.8%
2007 Q4	3,887	5,396	72.0%	191	280	68.2%	5,042	6,454	78.1%	9,120	12,130	75.2%
2008 Q1	4,347	6,020	72.2%	215	312	68.9%	5,624	7,142	78.7%	10,186	13,474	75.6%
2008 Q2	4,258	6,075	70.1%	218	315	69.2%	5,583	7,111	78.5%	10,059	13,501	74.5%
2008 Q3	4,302	6,108	70.4%	289	353	81.9%	5,538	6,961	79.6%	10,129	13,422	75.5%
2008 Q4	4,146	5,656	73.3%	247	336	73.5%	5,702	7,184	79.4%	10,095	13,176	76.6%
2009 Q1	4,791	6,467	74.1%	266	352	75.6%	5,884	7,633	77.1%	10,941	14,452	75.7%
2009 Q2	4,440	5,983	74.2%	239	320	74.7%	5,417	7,345	73.8%	10,096	13,648	74.0%
2009 Q3	4,551	6,131	74.2%	247	326	75.8%	5,517	7,386	74.7%	10,315	13,843	74.5%
2009 Q4	4,328	5,795	74.7%	195	259	75.3%	4,961	6,670	74.4%	9,484	12,724	74.5%
2010 Q1	4,883	6,402	76.3%	283	371	76.3%	5,774	7,636	75.6%	10,940	14,409	75.9%
2010 Q2	4,835	6,290	76.9%	230	314	73.2%	5,683	7,561	75.2%	10,748	14,165	75.9%
2010 Q3	4,902	6,309	77.7%	282	354	79.7%	5,745	7,596	75.6%	10,929	14,259	76.6%
2010 Q4	4,701	6,143	76.5%	237	320	74.1%	5,790	7,500	77.2%	10,728	13,963	76.8%
2011 Q1	5,144	6,896	74.6%	272	352	77.3%	6,212	8,120	76.5%	11,628	15,368	75.7%
2011 Q2	4,592	6,663	68.9%	286	363	78.8%	5,627	7,518	74.8%	10,505	14,544	72.2%
2011 Q3	5,374	7,529	71.4%	296	373	79.4%	6,047	7,968	75.9%	11,717	15,870	73.8%
2011 Q4	5,183	7,172	72.3%	309	394	78.4%	6,052	7,921	76.4%	11,544	15,487	74.5%
2012 Q1	6,085	8,331	73.0%	350	441	79.4%	6,597	8,582	76.9%	13,032	17,354	75.1%
2012 Q2	5,940	8,211	72.3%	369	450	82.0%	6,121	8,096	75.6%	12,430	16,757	74.2%

Table 6a.3 Proportion of individuals tested for syphilis who are tested for HIV at the same clinical encounter by HSDA – Females

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	Syphilis & HIV	Syphilis Test	%	Syphilis & HIV	Syphilis Test	%	Syphilis & HIV	Syphilis Test	%	Syphilis & HIV	Syphilis Test	%
2006 Q2	4,856	5,954	81.6%	563	684	82.3%	11,431	13,607	84.0%	16,850	20,245	83.2%
2006 Q3	5,237	6,268	83.6%	578	721	80.2%	12,182	14,369	84.8%	17,997	21,358	84.3%
2006 Q4	4,916	5,853	84.0%	549	652	84.2%	12,150	14,129	86.0%	17,615	20,634	85.4%
2007 Q1	5,505	6,423	85.7%	715	820	87.2%	14,139	16,268	86.9%	20,359	23,511	86.6%
2007 Q2	5,088	5,967	85.3%	623	725	85.9%	12,618	14,544	86.8%	18,329	21,236	86.3%
2007 Q3	5,291	6,041	87.6%	642	732	87.7%	12,909	14,843	87.0%	18,842	21,616	87.2%
2007 Q4	5,515	6,306	87.5%	653	749	87.2%	13,287	15,119	87.9%	19,455	22,174	87.7%
2008 Q1	6,108	7,026	86.9%	658	745	88.3%	14,497	16,362	88.6%	21,263	24,133	88.1%
2008 Q2	5,986	6,991	85.6%	672	778	86.4%	14,008	15,920	88.0%	20,666	23,689	87.2%
2008 Q3	5,903	6,799	86.8%	689	779	88.4%	14,055	15,956	88.1%	20,647	23,534	87.7%
2008 Q4	5,758	6,603	87.2%	642	723	88.8%	13,861	15,587	88.9%	20,261	22,913	88.4%
2009 Q1	6,257	7,020	89.1%	719	826	87.0%	15,297	17,357	88.1%	22,273	25,203	88.4%
2009 Q2	5,889	6,687	88.1%	651	751	86.7%	13,920	16,165	86.1%	20,460	23,603	86.7%
2009 Q3	5,986	6,787	88.2%	632	732	86.3%	14,188	16,460	86.2%	20,806	23,979	86.8%
2009 Q4	5,668	6,436	88.1%	664	742	89.5%	13,548	15,627	86.7%	19,880	22,805	87.2%
2010 Q1	6,347	7,144	88.8%	649	728	89.1%	15,177	17,437	87.0%	22,173	25,309	87.6%
2010 Q2	6,124	6,912	88.6%	602	688	87.5%	14,062	16,173	86.9%	20,788	23,773	87.4%
2010 Q3	6,323	7,023	90.0%	658	748	88.0%	14,614	16,742	87.3%	21,595	24,513	88.1%
2010 Q4	6,155	6,959	88.4%	605	705	85.8%	14,571	16,675	87.4%	21,331	24,339	87.6%
2011 Q1	6,667	7,566	88.1%	770	863	89.2%	15,394	17,602	87.5%	22,831	26,031	87.7%
2011 Q2	5,998	7,130	84.1%	629	701	89.7%	13,904	16,192	85.9%	20,531	24,023	85.5%
2011 Q3	6,610	7,650	86.4%	677	771	87.8%	14,765	17,198	85.9%	22,052	25,619	86.1%
2011 Q4	6,982	7,987	87.4%	727	828	87.8%	14,597	16,803	86.9%	22,306	25,618	87.1%
2012 Q1	8,222	9,338	88.0%	827	941	87.9%	16,019	18,411	87.0%	25,068	28,690	87.4%
2012 Q2	7,945	9,079	87.5%	720	825	87.3%	15,077	17,255	87.4%	23,742	27,159	87.4%

Table 6b.1 Proportion of individuals with a new STI diagnosis (e.g., syphilis) who are simultaneously tested for HIV by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	STI Dx & HIV Test	STI Dx	%	STI Dx & HIV Test	STI Dx	%	STI Dx & HIV Test	STI Dx	%	STI Dx & HIV Test	STI Dx	%
2006 Q2	184	559	32.9%	19	112	17.0%	436	1818	24.0%	639	2489	25.7%
2006 Q3	184	504	36.5%	31	127	24.4%	451	1914	23.6%	666	2545	26.2%
2006 Q4	187	538	34.8%	13	133	9.8%	392	1854	21.1%	592	2525	23.4%
2007 Q1	187	618	30.3%	26	145	17.9%	473	2084	22.7%	686	2847	24.1%
2007 Q2	185	532	34.8%	43	168	25.6%	400	1913	20.9%	628	2613	24.0%
2007 Q3	217	589	36.8%	25	151	16.6%	481	2065	23.3%	723	2805	25.8%
2007 Q4	196	561	34.9%	33	162	20.4%	513	1973	26.0%	742	2696	27.5%
2008 Q1	185	536	34.5%	39	168	23.2%	502	2113	23.8%	726	2817	25.8%
2008 Q2	217	600	36.2%	41	189	21.7%	521	2126	24.5%	779	2915	26.7%
2008 Q3	202	573	35.3%	32	162	19.8%	499	2232	22.4%	733	2967	24.7%
2008 Q4	217	577	37.6%	28	150	18.7%	582	2353	24.7%	827	3080	26.9%
2009 Q1	229	593	38.6%	37	158	23.4%	519	2146	24.2%	785	2897	27.1%
2009 Q2	195	548	35.6%	28	143	19.6%	505	2206	22.9%	728	2897	25.1%
2009 Q3	240	655	36.6%	31	180	17.2%	566	2412	23.5%	837	3247	25.8%
2009 Q4	228	589	38.7%	30	160	18.8%	491	2241	21.9%	749	2990	25.1%
2010 Q1	257	666	38.6%	32	150	21.3%	528	2438	21.7%	817	3254	25.1%
2010 Q2	232	626	37.1%	28	161	17.4%	513	2276	22.5%	773	3063	25.2%
2010 Q3	281	709	39.6%	37	187	19.8%	530	2264	23.4%	848	3160	26.8%
2010 Q4	311	750	41.5%	29	169	17.2%	550	2340	23.5%	890	3259	27.3%
2011 Q1	313	748	41.8%	28	180	15.6%	575	2321	24.8%	916	3249	28.2%
2011 Q2	298	739	40.3%	35	165	21.2%	530	2312	22.9%	863	3216	26.8%
2011 Q3	318	752	42.3%	27	154	17.5%	588	2400	24.5%	933	3306	28.2%
2011 Q4	300	716	41.9%	41	166	24.7%	567	2291	24.7%	908	3173	28.6%
2012 Q1	325	782	41.6%	40	167	24.0%	590	2316	25.5%	955	3265	29.2%

Table 6b.2 Proportion of individuals with a new STI diagnosis (e.g., syphilis) who are simultaneously tested for HIV HSDA – Males

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	STI Dx & HIV Test	STI Dx	%	STI Dx & HIV Test	STI Dx	%	STI Dx & HIV Test	STI Dx	%	STI Dx & HIV Test	STI Dx	%
2006 Q2	105	275	38.2%	6	35	17.1%	173	594	29.1%	284	904	31.4%
2006 Q3	95	233	40.8%	12	40	30.0%	192	651	29.5%	299	924	32.4%
2006 Q4	98	242	40.5%	7	45	15.6%	165	610	27.0%	270	897	30.1%
2007 Q1	100	282	35.5%	8	42	19.0%	208	717	29.0%	316	1041	30.4%
2007 Q2	93	241	38.6%	13	53	24.5%	156	663	23.5%	262	957	27.4%
2007 Q3	106	269	39.4%	9	52	17.3%	221	698	31.7%	336	1019	33.0%
2007 Q4	104	254	40.9%	18	65	27.7%	208	679	30.6%	330	998	33.1%
2008 Q1	105	245	42.9%	17	68	25.0%	216	742	29.1%	338	1055	32.0%
2008 Q2	123	262	46.9%	9	57	15.8%	212	725	29.2%	344	1044	33.0%
2008 Q3	99	253	39.1%	15	64	23.4%	227	805	28.2%	341	1122	30.4%
2008 Q4	108	252	42.9%	6	41	14.6%	255	867	29.4%	369	1160	31.8%
2009 Q1	130	265	49.1%	11	52	21.2%	241	781	30.9%	382	1098	34.8%
2009 Q2	107	242	44.2%	8	50	16.0%	219	754	29.0%	334	1046	31.9%
2009 Q3	124	283	43.8%	15	65	23.1%	233	824	28.3%	372	1172	31.7%
2009 Q4	131	287	45.6%	9	54	16.7%	200	781	25.6%	340	1122	30.3%
2010 Q1	137	297	46.1%	9	42	21.4%	219	823	26.6%	365	1162	31.4%
2010 Q2	111	250	44.4%	10	54	18.5%	239	763	31.3%	360	1067	33.7%
2010 Q3	160	346	46.2%	17	67	25.4%	228	751	30.4%	405	1164	34.8%
2010 Q4	185	354	52.3%	10	61	16.4%	239	806	29.7%	434	1221	35.5%
2011 Q1	181	360	50.3%	15	64	23.4%	240	792	30.3%	436	1216	35.9%
2011 Q2	158	338	46.7%	20	62	32.3%	208	745	27.9%	386	1145	33.7%
2011 Q3	177	394	44.9%	9	55	16.4%	254	816	31.1%	440	1265	34.8%
2011 Q4	175	383	45.7%	17	68	25.0%	236	803	29.4%	428	1254	34.1%
2012 Q1	204	417	48.9%	12	49	24.5%	257	830	31.0%	473	1296	36.5%

Table 6b.3 Proportion of individuals with a new STI diagnosis (e.g., syphilis) who are simultaneously tested for HIV HSDA – Females

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	STI Dx & HIV Test	STI Dx	%	STI Dx & HIV Test	STI Dx	%	STI Dx & HIV Test	STI Dx	%	STI Dx & HIV Test	STI Dx	%
2006 Q2	79	284	27.8%	13	77	16.9%	263	1223	21.5%	355	1584	22.4%
2006 Q3	89	271	32.8%	19	87	21.8%	259	1263	20.5%	367	1621	22.6%
2006 Q4	89	296	30.1%	6	88	6.8%	227	1243	18.3%	322	1627	19.8%
2007 Q1	87	336	25.9%	18	103	17.5%	265	1367	19.4%	370	1806	20.5%
2007 Q2	92	291	31.6%	30	115	26.1%	243	1248	19.5%	365	1654	22.1%
2007 Q3	111	319	34.8%	16	99	16.2%	260	1367	19.0%	387	1785	21.7%
2007 Q4	92	307	30.0%	15	97	15.5%	305	1294	23.6%	412	1698	24.3%
2008 Q1	80	290	27.6%	22	100	22.0%	286	1369	20.9%	388	1759	22.1%
2008 Q2	94	338	27.8%	32	132	24.2%	309	1400	22.1%	435	1870	23.3%
2008 Q3	103	320	32.2%	17	98	17.3%	272	1426	19.1%	392	1844	21.3%
2008 Q4	109	325	33.5%	22	108	20.4%	327	1485	22.0%	458	1918	23.9%
2009 Q1	99	328	30.2%	26	106	24.5%	278	1365	20.4%	403	1799	22.4%
2009 Q2	87	305	28.5%	20	93	21.5%	286	1451	19.7%	393	1849	21.3%
2009 Q3	114	369	30.9%	16	115	13.9%	333	1587	21.0%	463	2071	22.4%
2009 Q4	97	301	32.2%	21	106	19.8%	291	1460	19.9%	409	1867	21.9%
2010 Q1	120	369	32.5%	23	107	21.5%	309	1613	19.2%	452	2089	21.6%
2010 Q2	120	374	32.1%	18	107	16.8%	274	1513	18.1%	412	1994	20.7%
2010 Q3	121	363	33.3%	20	120	16.7%	302	1513	20.0%	443	1996	22.2%
2010 Q4	126	395	31.9%	19	108	17.6%	311	1534	20.3%	456	2037	22.4%
2011 Q1	132	388	34.0%	13	116	11.2%	335	1528	21.9%	480	2032	23.6%
2011 Q2	140	400	35.0%	15	103	14.6%	322	1567	20.5%	477	2070	23.0%
2011 Q3	140	357	39.2%	18	99	18.2%	334	1584	21.1%	492	2040	24.1%
2011 Q4	124	331	37.5%	24	98	24.5%	331	1487	22.3%	479	1916	25.0%
2012 Q1	121	365	33.2%	28	118	23.7%	333	1483	22.5%	482	1966	24.5%

Table 7.1 Proportion of individuals with a new HCV diagnosis who are tested for HIV within 3 months of HCV diagnosis by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	HCV +ve & HIV Test	HCV +	%	HCV +ve & HIV Test	HCV +	%	HCV +ve & HIV Test	HCV +	%	HCV +ve & HIV Test	HCV +	%
2006 Q3&4	194	371	52.3%	40	73	54.8%	553	1,047	52.8%	787	1,491	52.8%
2007 Q1&2	187	335	55.8%	41	65	63.1%	541	1,109	48.8%	769	1,509	51.0%
2007 Q3&4	190	323	58.8%	41	60	68.3%	508	1,000	50.8%	739	1,383	53.4%
2008 Q1&2	172	314	54.8%	42	63	66.7%	551	1,087	50.7%	765	1,464	52.3%
2008 Q3&4	181	295	61.4%	54	77	70.1%	456	939	48.6%	691	1,311	52.7%
2009 Q1&2	188	330	57.0%	41	71	57.7%	443	1,014	43.7%	672	1,415	47.5%
2009 Q3&4	154	287	53.7%	25	52	48.1%	363	831	43.7%	542	1,170	46.3%
2010 Q1&2	142	253	56.1%	27	43	62.8%	401	946	42.4%	570	1,242	45.9%
2010 Q3&4	132	234	56.4%	30	45	66.7%	351	809	43.4%	513	1,088	47.2%
2011 Q1&2	134	247	54.3%	28	45	62.2%	350	773	45.3%	512	1,065	48.1%
2011 Q3&4	132	242	54.5%	27	47	57.4%	325	750	43.3%	484	1,039	46.6%

Table 7.2 Proportion of individuals with a new HCV diagnosis who are tested for HIV within 3 months of HCV diagnosis by HSDA – Males

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	HCV +ve & HIV Test	HCV +	%	HCV +ve & HIV Test	HCV +	%	HCV +ve & HIV Test	HCV +	%	HCV +ve & HIV Test	HCV +	%
2006 Q3&4	126	235	53.6%	24	42	57.1%	354	678	52.2%	504	955	52.8%
2007 Q1&2	131	230	57.0%	25	42	59.5%	359	724	49.6%	515	996	51.7%
2007 Q3&4	122	210	58.1%	30	37	81.1%	317	639	49.6%	469	886	52.9%
2008 Q1&2	104	192	54.2%	30	43	69.8%	352	702	50.1%	486	937	51.9%
2008 Q3&4	126	207	60.9%	35	51	68.6%	288	592	48.6%	449	850	52.8%
2009 Q1&2	124	216	57.4%	25	47	53.2%	287	654	43.9%	436	917	47.5%
2009 Q3&4	102	189	54.0%	16	32	50.0%	249	539	46.2%	367	760	48.3%
2010 Q1&2	92	155	59.4%	16	26	61.5%	258	620	41.6%	366	801	45.7%
2010 Q3&4	87	153	56.9%	18	28	64.3%	248	557	44.5%	353	738	47.8%
2011 Q1&2	96	174	55.2%	19	32	59.4%	236	511	46.2%	351	717	49.0%
2011 Q3&4	86	155	55.5%	18	31	58.1%	223	499	44.7%	327	685	47.7%

Table 7.3 Proportion of individuals with a new HCV diagnosis who are tested for HIV within 3 months of HCV diagnosis by HSDA – Females

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	HCV +ve & HIV Test	HCV +	%	HCV +ve & HIV Test	HCV +	%	HCV +ve & HIV Test	HCV +	%	HCV +ve & HIV Test	HCV +	%
2006 Q3&4	67	135	49.6%	16	30	53.3%	191	356	53.7%	274	521	52.6%
2007 Q1&2	55	103	53.4%	16	21	76.2%	174	366	47.5%	245	490	50.0%
2007 Q3&4	68	112	60.7%	11	22	50.0%	181	346	52.3%	260	480	54.2%
2008 Q1&2	68	122	55.7%	12	20	60.0%	198	377	52.5%	278	519	53.6%
2008 Q3&4	55	88	62.5%	19	25	76.0%	167	344	48.5%	241	457	52.7%
2009 Q1&2	63	113	55.8%	15	23	65.2%	153	356	43.0%	231	492	47.0%
2009 Q3&4	52	97	53.6%	8	18	44.4%	112	288	38.9%	172	403	42.7%
2010 Q1&2	49	97	50.5%	11	17	64.7%	142	323	44.0%	202	437	46.2%
2010 Q3&4	45	81	55.6%	12	17	70.6%	103	250	41.2%	160	348	46.0%
2011 Q1&2	36	71	50.7%	9	13	69.2%	114	261	43.7%	159	345	46.1%
2011 Q3&4	45	85	52.9%	9	16	56.3%	102	250	40.8%	156	351	44.4%

Table 9.1 Proportion of individuals with a new HIV diagnosis with advanced HIV disease by HSDA

Year	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	HIV+ and AHD	HIV+	%	HIV+ and AHD	HIV+	%	HIV+ and AHD	HIV+	%	HIV+ and AHD	HIV+	%
2006	21	179	11.7%	2	19	10.5%	19	163	11.7%	42	361	11.6%
2007	14	192	7.3%	3	23	13.0%	23	176	13.1%	40	391	10.2%
2008	23	174	13.2%	0	10	~	29	161	18.0%	52	345	15.1%
2009	16	151	10.6%	1	16	6.3%	28	170	16.5%	45	337	13.4%
2010	12	148	8.1%	0	8	~	17	144	11.8%	29	300	9.7%

Table 9.2 Proportion of individuals with a new HIV diagnosis with advanced HIV disease by gender, BC

Year	Male			Female			Other			All		
	HIV+ and AHD	HIV+	%	HIV+ and AHD	HIV+	%	HIV+ and AHD	HIV+	%	HIV+ and AHD	HIV+	%
2006	37	289	12.8%	5	72	6.9%	0	0	---	42	361	11.6%
2007	36	303	11.9%	4	85	4.7%	0	3	---	40	391	10.2%
2008	43	286	15.0%	9	59	15.3%	0	0	---	52	345	15.1%
2009	41	266	15.4%	4	71	5.6%	0	0	---	45	337	13.4%
2010	24	238	10.0%	5	62	8.0%	0	2	---	29	300	10.0%

AHD = advanced HIV disease

Other = Transgender + Gender Unknown

Table 9.3 Proportion of individuals with a new HIV diagnosis with advanced HIV disease by Aboriginal status, BC

Year	Aboriginal			Non Aboriginal			Unknown			All		
	HIV+ and AHD	HIV+	%	HIV+ and AHD	HIV+	%	HIV+ and AHD	HIV+	%	HIV+ and AHD	HIV+	%
2006	6	55	11%	33	288	11%	3	18	17%	42	361	12%
2007	5	63	8%	31	307	10%	4	21	19%	40	391	10%
2008	5	46	11%	45	280	16%	2	19	11%	52	345	15%
2009	8	56	14%	36	265	14%	1	16	6%	45	337	13%
2010	2	39	5%	27	247	11%	0	14	0%	29	300	10%

AHD = advanced HIV disease

Unknown = Ethnicity unknown or not stated

Table 11.1 Proportion of individuals with a new HIV diagnosis with acute infection by HSDA

Year	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	HIV+ & Acute	HIV+	%	HIV+ & Acute	HIV+	%	HIV+ & Acute	HIV+	%	HIV+ & Acute	HIV+	%
2006	9	179	5%	1	19	5%	10	163	6%	20	361	6%
2007	18	192	9%	2	23	9%	5	176	3%	25	391	6%
2008	11	174	6%	0	10	0%	9	161	6%	20	345	6%
2009	13	151	9%	1	16	6%	5	170	3%	19	337	6%
2010	17	148	11%	0	8	0%	9	144	6%	26	300	9%
2011	16	163	10%	0	14	0%	7	112	6%	23	289	8%

Table 11.2 Proportion of individuals with a new HIV diagnosis with acute infection by gender, BC

Year	Male			Female			Other			All		
	HIV+ & Acute	HIV+	%	HIV+ & Acute	HIV+	%	HIV+ & Acute	HIV+	%	HIV+ & Acute	HIV+	%
2006	16	289	6%	4	72	6%	0	0	---	20	361	6%
2007	23	303	8%	2	85	2%	0	3	---	25	391	6%
2008	16	286	6%	4	59	7%	0	0	---	20	345	6%
2009	16	266	6%	3	71	4%	0	0	---	19	337	6%
2010	21	238	9%	5	62	8%	0	0	---	26	300	9%
2011	22	245	9%	1	43	2%	0	1	---	23	289	8%

Other = Transgender + Gender Unknown

Table 11.3 Proportion of individuals with a new HIV diagnosis with acute infection by Aboriginal status, BC

Year	Aboriginal			Non Aboriginal			Unknown		
	HIV+ & Acute	HIV+	%	HIV+ & Acute	HIV+	%	HIV+ & Acute	HIV+	%
2006	5	56	9%	14	288	5%	1	17	6%
2007	2	63	3%	23	307	7%	0	21	~
2008	2	46	4%	18	280	6%	0	19	~
2009	1	57	2%	17	264	6%	1	16	6%
2010	5	39	13%	20	248	8%	1	13	8%
2011	3	44	7%	20	228	9%	0	17	~

Unknown = Ethnicity unknown or not state

Table 14.1 Proportion of individuals starting antiretroviral therapy (ART) late in the course of HIV disease by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
2006 Q1	25 /	55	45.45%	6 /	6	100.00%	28 /	52	53.85%	59 /	113	52.21%
2006 Q2	29 /	53	54.72%	4 /	4	100.00%	20 /	41	48.78%	53 /	98	54.08%
2006 Q3	26 /	41	63.41%	0 /	1	0.00%	23 /	50	46.00%	49 /	92	53.26%
2006 Q4	26 /	48	54.17%	2 /	3	66.67%	30 /	58	51.72%	58 /	109	53.21%
2007 Q1	34 /	57	59.65%	5 /	6	83.33%	18 /	51	35.29%	57 /	114	50.00%
2007 Q2	30 /	73	41.10%	2 /	2	100.00%	35 /	61	57.38%	67 /	136	49.26%
2007 Q3	28 /	59	47.46%	0 /	1	0.00%	22 /	47	46.81%	50 /	107	46.73%
2007 Q4	23 /	58	39.66%	3 /	4	75.00%	27 /	61	44.26%	53 /	123	43.09%
2008 Q1	23 /	53	43.40%	0 /	0	0.00%	33 /	72	45.83%	56 /	125	44.80%
2008 Q2	20 /	58	34.48%	1 /	4	25.00%	31 /	67	46.27%	52 /	129	40.31%
2008 Q3	20 /	67	29.85%	3 /	6	50.00%	24 /	67	35.82%	47 /	140	33.57%
2008 Q4	19 /	56	33.93%	2 /	7	28.57%	22 /	68	32.35%	43 /	131	32.82%
2009 Q1	18 /	66	27.27%	3 /	7	42.86%	22 /	74	29.73%	43 /	147	29.25%
2009 Q2	15 /	58	25.86%	1 /	4	25.00%	28 /	79	35.44%	44 /	141	31.21%
2009 Q3	11 /	54	20.37%	3 /	6	50.00%	21 /	63	33.33%	35 /	123	28.46%
2009 Q4	16 /	58	27.59%	0 /	3	0.00%	19 /	82	23.17%	35 /	143	24.48%
2010 Q1	7 /	68	10.29%	0 /	0	0.00%	19 /	67	28.36%	26 /	135	19.26%
2010 Q2	14 /	59	23.73%	0 /	4	0.00%	23 /	62	37.10%	37 /	125	29.60%
2010 Q3	12 /	58	20.69%	2 /	3	66.67%	27 /	78	34.62%	41 /	139	29.50%
2010 Q4	16 /	54	29.63%	3 /	5	60.00%	19 /	61	31.15%	38 /	120	31.67%
2011 Q1	14 /	74	18.92%	7 /	9	77.78%	14 /	71	19.72%	35 /	154	22.73%
2011 Q2	13 /	71	18.31%	1 /	1	100.00%	15 /	62	24.19%	29 /	134	21.64%
2011 Q3	10 /	68	14.71%	2 /	8	25.00%	19 /	83	22.89%	31 /	159	19.50%
2011 Q4	7 /	58	12.07%	2 /	6	33.33%	20 /	69	28.99%	29 /	133	21.80%
2012 Q1	8 /	59	13.56%	1 /	3	33.33%	15 /	77	19.48%	24 /	139	17.27%
2012 Q2	8 /	44	18.18%	0 /	1	0.00%	13 /	74	17.57%	21 /	119	17.65%

Table 14.2 Proportion of individuals starting antiretroviral therapy (ART) late in the course of HIV disease by gender

Quarter	Male			Female		
	Count	Total	Percentage	Count	Total	Percentage
2006 Q1	50	95	52.63%	9	18	50.00%
2006 Q2	45	84	53.57%	8	14	57.14%
2006 Q3	35	70	50.00%	14	22	63.64%
2006 Q4	45	86	52.33%	13	23	56.52%
2007 Q1	43	87	49.43%	14	27	51.85%
2007 Q2	57	118	48.31%	10	18	55.56%
2007 Q3	40	82	48.78%	10	25	40.00%
2007 Q4	44	105	41.90%	9	18	50.00%
2008 Q1	45	103	43.69%	11	22	50.00%
2008 Q2	45	108	41.67%	7	21	33.33%
2008 Q3	31	106	29.25%	16	34	47.06%
2008 Q4	34	109	31.19%	9	22	40.91%
2009 Q1	34	116	29.31%	9	31	29.03%
2009 Q2	39	120	32.50%	5	21	23.81%
2009 Q3	31	104	29.81%	4	19	21.05%
2009 Q4	29	115	25.22%	6	28	21.43%
2010 Q1	18	107	16.82%	8	28	28.57%
2010 Q2	29	106	27.36%	8	19	42.11%
2010 Q3	31	110	28.18%	10	29	34.48%
2010 Q4	29	98	29.59%	9	22	40.91%
2011 Q1	28	128	21.88%	7	26	26.92%
2011 Q2	24	111	21.62%	5	23	21.74%
2011 Q3	24	130	18.46%	7	29	24.14%
2011 Q4	23	109	21.10%	6	24	25.00%
2012 Q1	21	113	18.58%	3	26	11.54%
2012 Q2	18	99	18.18%	3	20	15.00%

Table 18.1 Proportion of individuals with a new HIV diagnosis who are tested for syphilis within 3 months of HIV diagnosis by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%
2006 Q3&4	36	90	40%	3	9	33%	17	73	23%	56	172	33%
2007 Q1&2	66	124	53%	3	10	30%	21	76	28%	90	210	43%
2007 Q3&4	50	92	54%	5	14	36%	19	73	26%	74	179	41%
2008 Q1&2	75	111	68%	3	5	60%	21	63	33%	99	179	55%
2008 Q3&4	57	89	64%	3	7	43%	27	69	39%	87	165	53%
2009 Q1&2	58	96	60%	4	8	50%	33	81	41%	95	185	51%
2009 Q3&4	49	80	61%	3	10	30%	22	61	36%	74	151	49%
2010 Q1&2	59	88	67%	0	1	0%	27	65	42%	86	154	56%
2010 Q3&4	58	83	70%	2	7	29%	18	56	32%	78	146	53%
2011 Q1&2	51	76	67%	3	5	60%	22	57	39%	76	138	55%
2011 Q3&4	77	105	73%	3	9	33%	15	45	33%	95	159	60%

Table 18.2 Proportion of individuals with a new HIV diagnosis who are tested for syphilis within 3 months of HIV diagnosis – Males

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%
2006 Q3&4	32	71	45%	2	6	33%	14	58	24%	48	135	36%
2007 Q1&2	60	107	56%	1	6	17%	14	52	27%	75	165	45%
2007 Q3&4	41	74	55%	3	7	43%	14	55	25%	58	136	43%
2008 Q1&2	67	94	71%	0	1	0%	18	53	34%	85	148	57%
2008 Q3&4	53	78	68%	2	6	33%	18	52	35%	73	136	54%
2009 Q1&2	50	80	63%	2	4	50%	27	65	42%	79	149	53%
2009 Q3&4	43	70	61%	1	4	25%	14	43	33%	58	117	50%
2010 Q1&2	51	72	71%	0	1	0%	21	44	48%	72	117	62%
2010 Q3&4	52	72	72%	2	6	33%	17	43	40%	71	121	59%
2011 Q1&2	47	66	71%	2	3	67%	15	43	35%	64	112	57%
2011 Q3&4	73	96	76%	2	6	33%	13	35	37%	88	137	64%

Table 18.3 Proportion of individuals with a new HIV diagnosis who are tested for syphilis within 3 months of HIV diagnosis – Females

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%
2006 Q3&4	4	19	21%	1	3	33%	3	15	20%	8	37	22%
2007 Q1&2	6	16	38%	2	4	50%	6	22	27%	14	42	33%
2007 Q3&4	9	18	50%	2	7	29%	5	18	28%	16	43	37%
2008 Q1&2	8	17	47%	3	4	75%	3	10	30%	14	31	45%
2008 Q3&4	4	11	36%	1	1	100%	9	17	53%	14	29	48%
2009 Q1&2	8	16	50%	2	4	50%	6	16	38%	16	36	44%
2009 Q3&4	6	10	60%	2	6	33%	8	18	44%	16	34	47%
2010 Q1&2	8	16	50%	0	0	~	6	21	29%	14	37	38%
2010 Q3&4	6	11	55%	0	1	0%	1	13	8%	7	25	28%
2011 Q1&2	4	10	40%	1	2	50%	6	13	46%	11	25	44%
2011 Q3&4	4	8	50%	1	3	33%	2	10	20%	7	21	33%

Table 18.4 Proportion of individuals with a new HIV diagnosis who are tested for syphilis within 3 months of HIV diagnosis – by Aboriginal status

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%	HIV +ve & Syphilis test	HIV +ve	%
2006 Q3&4	3	9	33%	3	7	43%	4	11	36%	10	27	37%
2007 Q1&2	7	18	39%	3	8	38%	2	10	20%	12	36	33%
2007 Q3&4	3	7	43%	4	11	36%	3	9	33%	10	27	37%
2008 Q1&2	5	13	38%	3	4	75%	4	6	67%	12	23	52%
2008 Q3&4	7	10	70%	3	5	60%	4	8	50%	14	23	61%
2009 Q1&2	7	11	64%	2	2	100%	8	16	50%	17	29	59%
2009 Q3&4	4	9	44%	3	6	50%	6	12	50%	13	27	48%
2010 Q1&2	5	10	50%	0	0	n/a	3	10	30%	8	20	40%
2010 Q3&4	4	8	50%	1	4	n/a	1	7	14%	6	19	32%
2011 Q1&2	5	8	63%	1	2	n/a	1	7	14%	7	17	41%
2011 Q3&4	6	6	100%	1	1	n/a	1	3	33%	8	10	80%

Table 21.1 Percentage of HIV-infected individuals who are tested for genotypic antiretroviral drug resistance prior to starting antiretroviral therapy (ART) by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA		Other HSDAs			All HSDAs			
	Count	Count	Percentage	Count	Count	Percentage	Count	Count	Percentage	Count	Count	Percentage
2006 Q1	35	55	63.64%	5	6	83.33%	30	56	53.57%	70	117	59.83%
2006 Q2	38	53	71.70%	3	4	75.00%	20	45	44.44%	61	102	59.80%
2006 Q3	25	43	58.14%	1	1	100.00%	16	53	30.19%	42	97	43.30%
2006 Q4	36	48	75.00%	1	3	33.33%	38	60	63.33%	75	111	67.57%
2007 Q1	46	59	77.97%	5	6	83.33%	33	53	62.26%	84	118	71.19%
2007 Q2	58	75	77.33%	2	2	100.00%	37	64	57.81%	97	141	68.79%
2007 Q3	44	60	73.33%	0	1	0.00%	28	48	58.33%	72	109	66.06%
2007 Q4	45	59	76.27%	1	4	25.00%	47	68	69.12%	93	131	70.99%
2008 Q1	46	55	83.64%	0	0	0.00%	39	77	50.65%	85	132	64.39%
2008 Q2	41	58	70.69%	4	4	100.00%	47	71	66.20%	92	133	69.17%
2008 Q3	56	67	83.58%	4	6	66.67%	40	70	57.14%	100	143	69.93%
2008 Q4	39	58	67.24%	6	7	85.71%	52	70	74.29%	97	135	71.85%
2009 Q1	57	67	85.07%	6	7	85.71%	61	78	78.21%	124	152	81.58%
2009 Q2	47	58	81.03%	3	4	75.00%	64	82	78.05%	114	144	79.17%
2009 Q3	48	57	84.21%	6	6	100.00%	52	66	78.79%	106	129	82.17%
2009 Q4	50	59	84.75%	2	3	66.67%	65	83	78.31%	117	145	80.69%
2010 Q1	55	69	79.71%	0	0	0.00%	57	71	80.28%	112	140	80.00%
2010 Q2	53	61	86.89%	4	4	100.00%	55	66	83.33%	112	131	85.50%
2010 Q3	51	59	86.44%	3	3	100.00%	60	80	75.00%	114	142	80.28%
2010 Q4	44	55	80.00%	5	5	100.00%	49	65	75.38%	98	125	78.40%
2011 Q1	68	74	91.89%	9	9	100.00%	60	73	82.19%	137	156	87.82%
2011 Q2	63	73	86.30%	1	1	100.00%	52	66	78.79%	116	140	82.86%
2011 Q3	63	71	88.73%	8	8	100.00%	73	85	85.88%	144	164	87.80%
2011 Q4	49	58	84.48%	6	6	100.00%	57	73	78.08%	112	137	81.75%
2012 Q1	53	60	88.33%	3	4	75.00%	69	82	84.15%	125	146	85.62%
2012 Q2	40	45	88.89%	1	1	100.00%	64	81	79.01%	105	127	82.68%

Table 21.2 Percentage of HIV-infected individuals who are tested for genotypic antiretroviral drug resistance prior to starting antiretroviral therapy (ART) by gender

Quarter	Male			Female		
	Count	Total	Percentage	Count	Total	Percentage
2006 Q1	61	99	61.62%	9	18	50.00%
2006 Q2	54	86	62.79%	7	16	43.75%
2006 Q3	30	73	41.10%	12	24	50.00%
2006 Q4	62	88	70.45%	13	23	56.52%
2007 Q1	68	89	76.40%	16	29	55.17%
2007 Q2	87	122	71.31%	10	19	52.63%
2007 Q3	56	84	66.67%	16	25	64.00%
2007 Q4	81	112	72.32%	12	19	63.16%
2008 Q1	66	108	61.11%	19	24	79.17%
2008 Q2	78	111	70.27%	14	22	63.64%
2008 Q3	77	108	71.30%	23	35	65.71%
2008 Q4	81	112	72.32%	16	23	69.57%
2009 Q1	96	120	80.00%	28	32	87.50%
2009 Q2	98	123	79.67%	16	21	76.19%
2009 Q3	88	109	80.73%	18	20	90.00%
2009 Q4	95	117	81.20%	22	28	78.57%
2010 Q1	87	112	77.68%	25	28	89.29%
2010 Q2	95	110	86.36%	17	21	80.95%
2010 Q3	90	112	80.36%	24	30	80.00%
2010 Q4	78	101	77.23%	20	24	83.33%
2011 Q1	113	129	87.60%	24	27	88.89%
2011 Q2	95	114	83.33%	21	26	80.77%
2011 Q3	116	134	86.57%	28	30	93.33%
2011 Q4	91	113	80.53%	21	24	87.50%
2012 Q1	102	119	85.71%	23	27	85.19%
2012 Q2	88	107	82.24%	17	20	85.00%

Table 22.1 Percentage of individuals starting ART who achieve HIV plasma viral load (pVL) of < 200 copies/mL within 9 months of therapy initiation by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs						
2006 Q1	34	/	41	82.93%	1	/	1	100.00%	46	/	65	70.77%	81	/	107	75.70%
2006 Q2	41	/	53	77.36%	2	/	4	50.00%	28	/	43	65.12%	71	/	100	71.00%
2006 Q3	30	/	40	75.00%	1	/	2	50.00%	46	/	56	82.14%	77	/	98	78.57%
2006 Q4	43	/	55	78.18%	3	/	6	50.00%	43	/	56	76.79%	89	/	117	76.07%
2007 Q1	43	/	53	81.13%	2	/	4	50.00%	37	/	45	82.22%	82	/	102	80.39%
2007 Q2	33	/	43	76.74%	1	/	1	100.00%	39	/	53	73.58%	73	/	97	75.26%
2007 Q3	37	/	48	77.08%	0	/	3	0.00%	41	/	60	68.33%	78	/	111	70.27%
2007 Q4	48	/	59	81.36%	3	/	6	50.00%	39	/	53	73.58%	90	/	118	76.27%
2008 Q1	62	/	75	82.67%	1	/	2	50.00%	46	/	64	71.88%	109	/	141	77.30%
2008 Q2	53	/	60	88.33%	1	/	1	100.00%	39	/	48	81.25%	93	/	109	85.32%
2008 Q3	49	/	59	83.05%	4	/	4	100.00%	57	/	68	83.82%	110	/	131	83.97%
2008 Q4	45	/	55	81.82%	0	/	0	0.00%	63	/	77	81.82%	108	/	132	81.82%
2009 Q1	49	/	58	84.48%	2	/	4	50.00%	58	/	71	81.69%	109	/	133	81.95%
2009 Q2	60	/	67	89.55%	3	/	6	50.00%	59	/	70	84.29%	122	/	143	85.31%
2009 Q3	48	/	58	82.76%	3	/	7	42.86%	62	/	70	88.57%	113	/	135	83.70%
2009 Q4	59	/	67	88.06%	6	/	7	85.71%	66	/	78	84.62%	131	/	152	86.18%
2010 Q1	50	/	58	86.21%	2	/	4	50.00%	70	/	82	85.37%	122	/	144	84.72%
2010 Q2	51	/	57	89.47%	4	/	6	66.67%	52	/	66	78.79%	107	/	129	82.95%
2010 Q3	51	/	59	86.44%	3	/	3	100.00%	67	/	83	80.72%	121	/	145	83.45%
2010 Q4	61	/	69	88.41%	0	/	0	0.00%	60	/	71	84.51%	121	/	140	86.43%
2011 Q1	55	/	61	90.16%	3	/	4	75.00%	53	/	66	80.30%	111	/	131	84.73%
2011 Q2	53	/	59	89.83%	2	/	3	66.67%	74	/	80	92.50%	129	/	142	90.85%
2011 Q3	45	/	55	81.82%	4	/	5	80.00%	49	/	65	75.38%	98	/	125	78.40%
2011 Q4	64	/	74	86.49%	4	/	9	44.44%	58	/	73	79.45%	126	/	156	80.77%
2012 Q1	67	/	73	91.78%	1	/	1	100.00%	57	/	66	86.36%	125	/	140	89.29%
2012 Q2	65	/	71	91.55%	6	/	8	75.00%	71	/	85	83.53%	142	/	164	86.59%

Table 22.2 Percentage of individuals starting ART who achieve HIV plasma viral load (pVL) of < 50 copies/mL within 9 months of therapy initiation by gender

Quarter	Male			Female		
	Count	Total	Percentage	Count	Total	Percentage
2006 Q1	68	87	78.16%	13	20	65.00%
2006 Q2	57	80	71.25%	14	20	70.00%
2006 Q3	66	80	82.50%	11	18	61.11%
2006 Q4	77	99	77.78%	12	18	66.67%
2007 Q1	71	86	82.56%	11	16	68.75%
2007 Q2	55	73	75.34%	18	24	75.00%
2007 Q3	65	88	73.86%	13	23	56.52%
2007 Q4	68	89	76.40%	22	29	75.86%
2008 Q1	98	122	80.33%	11	19	57.89%
2008 Q2	72	84	85.71%	21	25	84.00%
2008 Q3	99	112	88.39%	11	19	57.89%
2008 Q4	93	108	86.11%	15	24	62.50%
2009 Q1	92	111	82.88%	17	22	77.27%
2009 Q2	94	108	87.04%	28	35	80.00%
2009 Q3	98	112	87.50%	15	23	65.22%
2009 Q4	103	120	85.83%	28	32	87.50%
2010 Q1	106	123	86.18%	16	21	76.19%
2010 Q2	90	109	82.57%	17	20	85.00%
2010 Q3	96	117	82.05%	25	28	89.29%
2010 Q4	96	112	85.71%	25	28	89.29%
2011 Q1	93	110	84.55%	18	21	85.71%
2011 Q2	102	112	91.07%	27	30	90.00%
2011 Q3	81	101	80.20%	17	24	70.83%
2011 Q4	105	129	81.40%	21	27	77.78%
2012 Q1	104	114	91.23%	21	26	80.77%
2012 Q2	124	134	92.54%	18	30	60.00%

Table 23.1 Percentage of individuals who initiated antiretroviral therapy (ART) with a recommended therapy regimen (among those with no drug resistance) by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs						
2006 Q1	19	/	27	70.37%	3	/	3	100.00%	17	/	21	80.95%	39	/	51	76.47%
2006 Q2	20	/	30	66.67%	2	/	2	100.00%	14	/	17	82.35%	36	/	49	73.47%
2006 Q3	15	/	22	68.18%	1	/	1	100.00%	11	/	14	78.57%	27	/	37	72.97%
2006 Q4	23	/	30	76.67%	0	/	1	0.00%	22	/	35	62.86%	45	/	66	68.18%
2007 Q1	34	/	44	77.27%	5	/	5	100.00%	23	/	32	71.88%	62	/	81	76.54%
2007 Q2	48	/	55	87.27%	2	/	2	100.00%	26	/	35	74.29%	76	/	92	82.61%
2007 Q3	31	/	38	81.58%	0	/	0	0.00%	24	/	27	88.89%	55	/	65	84.62%
2007 Q4	34	/	43	79.07%	1	/	1	100.00%	25	/	37	67.57%	60	/	81	74.07%
2008 Q1	30	/	41	73.17%	0	/	0	0.00%	29	/	36	80.56%	59	/	77	76.62%
2008 Q2	30	/	38	78.95%	2	/	2	100.00%	25	/	39	64.10%	57	/	79	72.15%
2008 Q3	43	/	50	86.00%	4	/	4	100.00%	29	/	36	80.56%	76	/	90	84.44%
2008 Q4	32	/	36	88.89%	4	/	4	100.00%	39	/	46	84.78%	75	/	86	87.21%
2009 Q1	46	/	52	88.46%	6	/	6	100.00%	49	/	56	87.50%	101	/	114	88.60%
2009 Q2	33	/	39	84.62%	1	/	3	33.33%	51	/	58	87.93%	85	/	100	85.00%
2009 Q3	38	/	46	82.61%	6	/	6	100.00%	42	/	49	85.71%	86	/	101	85.15%
2009 Q4	39	/	43	90.70%	2	/	2	100.00%	52	/	57	91.23%	93	/	102	91.18%
2010 Q1	41	/	52	78.85%	0	/	0	0.00%	43	/	49	87.76%	84	/	101	83.17%
2010 Q2	38	/	46	82.61%	4	/	4	100.00%	40	/	51	78.43%	82	/	101	81.19%
2010 Q3	37	/	47	78.72%	3	/	3	100.00%	40	/	53	75.47%	80	/	103	77.67%
2010 Q4	30	/	40	75.00%	3	/	3	100.00%	38	/	45	84.44%	71	/	88	80.68%
2011 Q1	56	/	65	86.15%	7	/	9	77.78%	48	/	55	87.27%	111	/	129	86.05%
2011 Q2	39	/	54	72.22%	1	/	1	100.00%	35	/	43	81.40%	75	/	98	76.53%
2011 Q3	46	/	57	80.70%	8	/	8	100.00%	53	/	66	80.30%	107	/	131	81.68%
2011 Q4	28	/	40	70.00%	6	/	6	100.00%	34	/	51	66.67%	68	/	97	70.10%
2012 Q1	37	/	47	78.72%	2	/	3	66.67%	45	/	59	76.27%	84	/	109	77.06%
2012 Q2	26	/	36	72.22%	1	/	1	100.00%	43	/	57	75.44%	70	/	94	74.47%

Table 23.2 Percentage of individuals who initiated antiretroviral therapy (ART) with a recommended therapy regimen (among those with no drug resistance) by gender

Quarter	Male			Female		
	Count	Denominator	Percentage	Count	Denominator	Percentage
2006 Q1	35	46	76.09%	4	5	80.00%
2006 Q2	33	45	73.33%	3	4	75.00%
2006 Q3	20	25	80.00%	7	12	58.33%
2006 Q4	38	54	70.37%	7	12	58.33%
2007 Q1	52	65	80.00%	10	16	62.50%
2007 Q2	70	82	85.37%	6	10	60.00%
2007 Q3	43	50	86.00%	12	15	80.00%
2007 Q4	51	70	72.86%	9	11	81.82%
2008 Q1	46	61	75.41%	13	16	81.25%
2008 Q2	49	65	75.38%	8	14	57.14%
2008 Q3	60	69	86.96%	16	21	76.19%
2008 Q4	62	71	87.32%	13	15	86.67%
2009 Q1	84	92	91.30%	17	22	77.27%
2009 Q2	76	85	89.41%	9	15	60.00%
2009 Q3	72	83	86.75%	14	18	77.78%
2009 Q4	74	82	90.24%	19	20	95.00%
2010 Q1	67	78	85.90%	17	23	73.91%
2010 Q2	74	85	87.06%	8	16	50.00%
2010 Q3	63	81	77.78%	17	22	77.27%
2010 Q4	58	71	81.69%	13	17	76.47%
2011 Q1	95	106	89.62%	16	23	69.57%
2011 Q2	60	79	75.95%	15	19	78.95%
2011 Q3	86	105	81.90%	21	26	80.77%
2011 Q4	57	80	71.25%	11	17	64.71%
2012 Q1	71	91	78.02%	13	18	72.22%
2012 Q2	61	80	76.25%	9	14	64.29%

Table 24.1 Percentage of individuals on antiretroviral therapy (ART) that achieve annual prescription refill adherence of > 95% by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs						
2006 Q1	1280	/	1781	71.87%	16	/	32	50.00%	994	/	1497	66.40%	2290	/	3310	69.18%
2006 Q2	1343	/	1825	73.59%	17	/	31	54.84%	1028	/	1554	66.15%	2388	/	3410	70.03%
2006 Q3	1389	/	1859	74.72%	17	/	31	54.84%	1089	/	1564	69.63%	2495	/	3454	72.24%
2006 Q4	1409	/	1874	75.19%	16	/	30	53.33%	1113	/	1628	68.37%	2538	/	3532	71.86%
2007 Q1	1478	/	1914	77.22%	20	/	36	55.56%	1134	/	1675	67.70%	2632	/	3625	72.61%
2007 Q2	1491	/	1992	74.85%	21	/	36	58.33%	1152	/	1730	66.59%	2664	/	3758	70.89%
2007 Q3	1533	/	2046	74.93%	19	/	33	57.58%	1174	/	1748	67.16%	2726	/	3827	71.23%
2007 Q4	1549	/	2089	74.15%	23	/	42	54.76%	1213	/	1760	68.92%	2785	/	3891	71.58%
2008 Q1	1596	/	2149	74.27%	20	/	43	46.51%	1236	/	1775	69.63%	2852	/	3967	71.89%
2008 Q2	1654	/	2232	74.10%	18	/	46	39.13%	1289	/	1856	69.45%	2961	/	4134	71.63%
2008 Q3	1664	/	2272	73.24%	18	/	49	36.73%	1321	/	1896	69.67%	3003	/	4217	71.21%
2008 Q4	1726	/	2332	74.01%	23	/	54	42.59%	1362	/	1941	70.17%	3111	/	4327	71.90%
2009 Q1	1762	/	2377	74.13%	23	/	54	42.59%	1421	/	2011	70.66%	3206	/	4442	72.17%
2009 Q2	1817	/	2436	74.59%	29	/	56	51.79%	1429	/	2078	68.77%	3275	/	4570	71.66%
2009 Q3	1885	/	2506	75.22%	33	/	58	56.90%	1487	/	2118	70.21%	3405	/	4682	72.73%
2009 Q4	1946	/	2537	76.70%	37	/	62	59.68%	1516	/	2170	69.86%	3499	/	4769	73.37%
2010 Q1	2000	/	2601	76.89%	36	/	67	53.73%	1549	/	2216	69.90%	3585	/	4884	73.40%
2010 Q2	2020	/	2657	76.03%	36	/	77	46.75%	1590	/	2255	70.51%	3646	/	4989	73.08%
2010 Q3	2054	/	2722	75.46%	40	/	78	51.28%	1618	/	2317	69.83%	3712	/	5117	72.54%
2010 Q4	2065	/	2766	74.66%	41	/	78	52.56%	1653	/	2379	69.48%	3759	/	5223	71.97%
2011 Q1	2144	/	2846	75.33%	43	/	81	53.09%	1667	/	2414	69.06%	3854	/	5341	72.16%
2011 Q2	2180	/	2879	75.72%	45	/	88	51.14%	1687	/	2461	68.55%	3912	/	5428	72.07%
2011 Q3	2214	/	2946	75.15%	42	/	92	45.65%	1718	/	2500	68.72%	3974	/	5538	71.76%
2011 Q4	2280	/	2988	76.31%	49	/	102	48.04%	1719	/	2535	67.81%	4048	/	5625	71.96%
2012 Q1	2349	/	3041	77.24%	55	/	103	53.40%	1769	/	2598	68.09%	4173	/	5742	72.68%
2012 Q2	2368	/	3059	77.41%	64	/	108	59.26%	1833	/	2665	68.78%	4265	/	5832	73.13%

Table 24.2 Percentage of individuals on antiretroviral therapy (ART) that achieve annual prescription refill adherence of > 95% by gender

Quarter	Male				Female			
	2024	/	2853	70.94%	266	/	457	58.21%
2006 Q2	2120	/	2928	72.40%	268	/	482	55.60%
2006 Q3	2209	/	2961	74.60%	286	/	493	58.01%
2006 Q4	2241	/	3029	73.98%	297	/	503	59.05%
2007 Q1	2329	/	3112	74.84%	303	/	513	59.06%
2007 Q2	2365	/	3235	73.11%	299	/	523	57.17%
2007 Q3	2420	/	3285	73.67%	306	/	542	56.46%
2007 Q4	2472	/	3338	74.06%	313	/	553	56.60%
2008 Q1	2516	/	3389	74.24%	336	/	578	58.13%
2008 Q2	2627	/	3534	74.34%	334	/	600	55.67%
2008 Q3	2644	/	3585	73.75%	359	/	632	56.80%
2008 Q4	2741	/	3688	74.32%	370	/	639	57.90%
2009 Q1	2819	/	3770	74.77%	387	/	672	57.59%
2009 Q2	2869	/	3868	74.17%	406	/	702	57.83%
2009 Q3	2973	/	3963	75.02%	432	/	719	60.08%
2009 Q4	3056	/	4030	75.83%	443	/	739	59.95%
2010 Q1	3126	/	4121	75.86%	459	/	763	60.16%
2010 Q2	3181	/	4210	75.56%	465	/	779	59.69%
2010 Q3	3254	/	4325	75.24%	458	/	792	57.83%
2010 Q4	3296	/	4395	74.99%	463	/	828	55.92%
2011 Q1	3372	/	4485	75.18%	482	/	856	56.31%
2011 Q2	3420	/	4540	75.33%	492	/	888	55.41%
2011 Q3	3455	/	4644	74.40%	519	/	894	58.05%
2011 Q4	3506	/	4715	74.36%	542	/	910	59.56%
2012 Q1	3614	/	4807	75.18%	559	/	935	59.79%
2012 Q2	3687	/	4877	75.60%	578	/	955	60.52%

Table 25.1 Number of physicians initiating therapy or providing HIV-related care to patients on antiretroviral therapy (ART)

Quarter	Vancouver HSDA	Northern Interior HSDA	Other HSDAs	All HSDAs
2006 Q1	162	19	402	583
2006 Q2	165	22	407	594
2006 Q3	171	22	407	600
2006 Q4	160	22	417	599
2007 Q1	168	24	424	616
2007 Q2	166	22	455	643
2007 Q3	168	21	438	627
2007 Q4	163	22	455	640
2008 Q1	174	22	442	638
2008 Q2	177	25	470	672
2008 Q3	178	28	464	670
2008 Q4	183	27	461	671
2009 Q1	191	27	489	707
2009 Q2	184	26	499	709
2009 Q3	191	24	482	697
2009 Q4	187	23	502	712
2010 Q1	185	23	498	706
2010 Q2	192	21	504	717
2010 Q3	192	23	512	727
2010 Q4	189	23	511	723
2011 Q1	196	25	521	742
2011 Q2	193	27	525	745
2011 Q3	200	26	528	754
2011 Q4	199	24	530	753
2012 Q1	199	25	517	741
2012 Q2	201	25	522	748

Table 26.1 Percentage of individuals on antiretroviral therapy (ART) who experience a serious adverse drug reaction (ADR) by HSDA

Quarter	Vancouver HSDA		Northern Interior HSDA		Other HSDAs		All HSDAs					
2006 Q1	/	1968	/	45	/	1686	/	3699				
2006 Q2	/	2005	/	46	/	1740	/	3791				
2006 Q3	/	2031	/	41	/	1765	/	3837				
2006 Q4	/	2073	/	42	/	1805	/	3920				
2007 Q1	/	2121	/	44	/	1842	/	4007				
2007 Q2	/	2217	/	50	/	1920	/	4187				
2007 Q3	/	2286	/	46	/	1931	/	4263				
2007 Q4	/	2342	/	53	/	1956	/	4351				
2008 Q1	1/	2382	0.04%	0/	52	0.00%	0/	2008	0.00%	1/	4442	0.02%
2008 Q2	1/	2457	0.04%	0/	55	0.00%	1/	2099	0.05%	2/	4611	0.04%
2008 Q3	5/	2517	0.20%	0/	58	0.00%	1/	2149	0.05%	6/	4724	0.13%
2008 Q4	1/	2587	0.04%	0/	66	0.00%	3/	2189	0.14%	4/	4842	0.08%
2009 Q1	5/	2657	0.19%	1/	76	1.32%	2/	2242	0.09%	8/	4975	0.16%
2009 Q2	6/	2718	0.22%	0/	80	0.00%	3/	2322	0.13%	9/	5120	0.18%
2009 Q3	1/	2783	0.04%	0/	79	0.00%	3/	2346	0.13%	4/	5208	0.08%
2009 Q4	2/	2825	0.07%	0/	81	0.00%	3/	2403	0.12%	5/	5309	0.09%
2010 Q1	3/	2875	0.10%	0/	81	0.00%	4/	2464	0.16%	7/	5420	0.13%
2010 Q2	2/	2915	0.07%	0/	89	0.00%	0/	2504	0.00%	2/	5508	0.04%
2010 Q3	1/	3009	0.03%	0/	86	0.00%	1/	2549	0.04%	2/	5644	0.04%
2010 Q4	4/	3039	0.13%	0/	93	0.00%	1/	2611	0.04%	5/	5743	0.09%
2011 Q1	2/	3118	0.06%	0/	105	0.00%	1/	2656	0.04%	3/	5879	0.05%
2011 Q2	1/	3171	0.03%	1/	106	0.94%	2/	2690	0.07%	4/	5967	0.07%
2011 Q3	2/	3220	0.06%	0/	112	0.00%	7/	2773	0.25%	9/	6105	0.15%
2011 Q4	2/	3241	0.06%	0/	122	0.00%	1/	2837	0.04%	3/	6200	0.05%
2012 Q1	3/	3293	0.09%	0/	118	0.00%	3/	2891	0.10%	6/	6302	0.10%
2012 Q2	3/	3299	0.09%	0/	122	0.00%	1/	2959	0.03%	4/	6380	0.06%

Table 26.2 Percentage of individuals on antiretroviral therapy (ART) who experience a serious adverse drug reaction (ADR) by gender

Quarter	Male			Female		
2006 Q1	/	3174		/	525	
2006 Q2	/	3248		/	543	
2006 Q3	/	3280		/	557	
2006 Q4	/	3351		/	569	
2007 Q1	/	3418		/	589	
2007 Q2	/	3580		/	607	
2007 Q3	/	3636		/	627	
2007 Q4	/	3723		/	628	
2008 Q1	1 /	3792	0.03%	0 /	650	0.00%
2008 Q2	1 /	3928	0.03%	1 /	683	0.15%
2008 Q3	6 /	4000	0.15%	0 /	724	0.00%
2008 Q4	3 /	4110	0.07%	1 /	732	0.14%
2009 Q1	5 /	4201	0.12%	3 /	774	0.39%
2009 Q2	6 /	4317	0.14%	3 /	803	0.37%
2009 Q3	4 /	4404	0.09%	0 /	804	0.00%
2009 Q4	4 /	4477	0.09%	1 /	832	0.12%
2010 Q1	4 /	4566	0.09%	3 /	854	0.35%
2010 Q2	1 /	4637	0.02%	1 /	871	0.11%
2010 Q3	1 /	4751	0.02%	1 /	893	0.11%
2010 Q4	3 /	4817	0.06%	2 /	926	0.22%
2011 Q1	3 /	4926	0.06%	0 /	953	0.00%
2011 Q2	2 /	4979	0.04%	2 /	988	0.20%
2011 Q3	6 /	5107	0.12%	3 /	998	0.30%
2011 Q4	2 /	5186	0.04%	1 /	1014	0.10%
2012 Q1	5 /	5269	0.09%	1 /	1033	0.10%
2012 Q2	3 /	5330	0.06%	1 /	1050	0.10%

Table 28.1 Incidence of resistance to any antiretroviral drug by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs		
2006 Q1	11 /	5406.2	0.20%	0 /	102	0.00%	8 /	4556	0.18%	19 /	10074	0.19%
2006 Q2	13 /	5551.7	0.23%	1 /	112	0.89%	5 /	4673	0.11%	19 /	10345	0.18%
2006 Q3	11 /	5608.8	0.20%	0 /	102	0.00%	10 /	4777	0.21%	21 /	10500	0.20%
2006 Q4	11 /	5750.4	0.19%	0 /	113	0.00%	7 /	4846	0.14%	18 /	10730	0.17%
2007 Q1	10 /	5893.9	0.17%	0 /	110	0.00%	8 /	5023	0.16%	18 /	11045	0.16%
2007 Q2	4 /	6052.9	0.07%	1 /	119	0.84%	4 /	5130	0.08%	10 /	11316	0.09%
2007 Q3	8 /	6360.9	0.13%	0 /	118	0.00%	4 /	5236	0.08%	12 /	11727	0.10%
2007 Q4	7 /	6490.4	0.11%	2 /	121	1.66%	5 /	5310	0.09%	14 /	11937	0.12%
2008 Q1	8 /	6622.4	0.12%	1 /	134	0.75%	7 /	5435	0.13%	16 /	12205	0.13%
2008 Q2	11 /	6757.5	0.16%	0 /	135	0.00%	8 /	5684	0.14%	19 /	12596	0.15%
2008 Q3	1 /	6899.4	0.01%	2 /	150	1.33%	5 /	5826	0.09%	8 /	12889	0.06%
2008 Q4	11 /	7205.2	0.15%	0 /	171	0.00%	2 /	6006	0.03%	13 /	13397	0.10%
2009 Q1	9 /	7365.9	0.12%	3 /	185	1.62%	10 /	6142	0.16%	22 /	13710	0.16%
2009 Q2	7 /	7637.6	0.09%	0 /	213	0.00%	5 /	6312	0.08%	12 /	14184	0.08%
2009 Q3	5 /	7797.8	0.06%	1 /	200	0.50%	6 /	6428	0.09%	12 /	14449	0.08%
2009 Q4	12 /	7882.7	0.15%	0 /	215	0.00%	7 /	6551	0.11%	20 /	14677	0.14%
2010 Q1	5 /	7960.8	0.06%	0 /	215	0.00%	11 /	6729	0.16%	16 /	14929	0.11%
2010 Q2	6 /	8145.9	0.07%	1 /	229	0.44%	4 /	6869	0.06%	11 /	15260	0.07%
2010 Q3	6 /	8406	0.07%	1 /	217	0.46%	7 /	6946	0.10%	14 /	15582	0.09%
2010 Q4	8 /	8458.1	0.09%	1 /	243	0.41%	8 /	7125	0.11%	17 /	15850	0.11%
2011 Q1	11 /	8691.8	0.13%	2 /	257	0.78%	6 /	7203	0.08%	19 /	16186	0.12%
2011 Q2	13 /	8871	0.15%	2 /	268	0.75%	11 /	7261	0.15%	26 /	16446	0.16%
2011 Q3	8 /	9032.5	0.09%	0 /	296	0.00%	8 /	7482	0.11%	16 /	16884	0.09%
2011 Q4	5 /	9146.1	0.05%	1 /	313	0.32%	6 /	7711	0.08%	12 /	17252	0.07%
2012 Q1	9 /	9282.1	0.10%	1 /	311	0.32%	10 /	7874	0.13%	20 /	17596	0.11%
2012 Q2	5 /	9264.4	0.05%	1 /	326	0.31%	6 /	7999	0.08%	13 /	17751	0.07%

Table 28.2 Incidence of resistance to any antiretroviral drug by gender (BC)

Quarter	Male			Female		
	Count	Rate	%	Count	Rate	%
2006 Q1	12	8714.6	0.14%	7	1360	0.51%
2006 Q2	9	8955.9	0.10%	10	1389	0.72%
2006 Q3	18	9069.6	0.20%	3	1430	0.21%
2006 Q4	12	9273	0.13%	6	1457	0.41%
2007 Q1	13	9537.9	0.14%	5	1507	0.33%
2007 Q2	9	9760.9	0.09%	1	1555	0.06%
2007 Q3	8	10100	0.08%	4	1626	0.25%
2007 Q4	10	10298	0.10%	4	1639	0.24%
2008 Q1	14	10524	0.13%	2	1680	0.12%
2008 Q2	15	10843	0.14%	4	1753	0.23%
2008 Q3	6	11056	0.05%	2	1833	0.11%
2008 Q4	12	11447	0.10%	1	1950	0.05%
2009 Q1	15	11680	0.13%	7	2030	0.34%
2009 Q2	8	12076	0.07%	4	2108	0.19%
2009 Q3	9	12307	0.07%	3	2142	0.14%
2009 Q4	17	12489	0.14%	3	2187	0.14%
2010 Q1	11	12708	0.09%	5	2221	0.23%
2010 Q2	6	12950	0.05%	5	2310	0.22%
2010 Q3	10	13263	0.08%	4	2319	0.17%
2010 Q4	13	13445	0.10%	4	2405	0.17%
2011 Q1	13	13676	0.10%	6	2510	0.24%
2011 Q2	13	13880	0.09%	13	2566	0.51%
2011 Q3	11	14213	0.08%	5	2672	0.19%
2011 Q4	4	14556	0.03%	8	2696	0.30%
2012 Q1	17	14834	0.11%	3	2761	0.11%
2012 Q2	9	14943	0.06%	4	2808	0.14%

Table 29.1 Proportion of individuals on antiretroviral therapy who change antiretroviral drug treatment by HSDA

Quarter	Vancouver HSDA			Northern Interior HSDA			Other HSDAs			All HSDAs						
2006 Q1	182	/	1968	9.25%	3	/	45	6.67%	138	/	1682	8.20%	323	/	3699	8.73%
2006 Q2	220	/	2005	10.97%	2	/	46	4.35%	183	/	1736	10.54%	406	/	3791	10.71%
2006 Q3	233	/	2031	11.47%	2	/	41	4.88%	181	/	1761	10.28%	416	/	3837	10.84%
2006 Q4	244	/	2073	11.77%	3	/	42	7.14%	209	/	1794	11.65%	458	/	3920	11.68%
2007 Q1	242	/	2121	11.41%	1	/	44	2.27%	165	/	1833	9.00%	408	/	4006	10.18%
2007 Q2	219	/	2217	9.88%	0	/	50	0.00%	168	/	1914	8.78%	388	/	4187	9.27%
2007 Q3	156	/	2286	6.82%	2	/	46	4.35%	120	/	1926	6.23%	279	/	4263	6.54%
2007 Q4	226	/	2342	9.65%	1	/	53	1.89%	166	/	1950	8.51%	393	/	4351	9.03%
2008 Q1	202	/	2382	8.48%	1	/	52	1.92%	163	/	1999	8.15%	366	/	4441	8.24%
2008 Q2	259	/	2457	10.54%	5	/	55	9.09%	193	/	2088	9.24%	458	/	4611	9.93%
2008 Q3	206	/	2516	8.19%	2	/	58	3.45%	134	/	2144	6.25%	342	/	4724	7.24%
2008 Q4	162	/	2587	6.26%	1	/	66	1.52%	153	/	2180	7.02%	316	/	4841	6.53%
2009 Q1	176	/	2657	6.62%	2	/	76	2.63%	146	/	2234	6.54%	324	/	4975	6.51%
2009 Q2	146	/	2718	5.37%	5	/	80	6.25%	117	/	2312	5.06%	269	/	5120	5.25%
2009 Q3	123	/	2784	4.42%	3	/	79	3.80%	100	/	2335	4.28%	226	/	5208	4.34%
2009 Q4	120	/	2825	4.25%	2	/	81	2.47%	105	/	2392	4.39%	227	/	5309	4.28%
2010 Q1	133	/	2875	4.63%	2	/	81	2.47%	99	/	2454	4.03%	234	/	5420	4.32%
2010 Q2	117	/	2915	4.01%	0	/	89	0.00%	101	/	2497	4.04%	218	/	5508	3.96%
2010 Q3	137	/	3008	4.55%	1	/	86	1.16%	112	/	2543	4.40%	250	/	5644	4.43%
2010 Q4	131	/	3039	4.31%	5	/	93	5.38%	110	/	2602	4.23%	246	/	5743	4.28%
2011 Q1	131	/	3119	4.20%	4	/	105	3.81%	119	/	2638	4.51%	255	/	5879	4.34%
2011 Q2	122	/	3171	3.85%	4	/	106	3.77%	93	/	2672	3.48%	219	/	5967	3.67%
2011 Q3	132	/	3219	4.10%	10	/	112	8.93%	105	/	2747	3.82%	247	/	6105	4.05%
2011 Q4	113	/	3241	3.49%	5	/	122	4.10%	82	/	2794	2.93%	200	/	6200	3.23%
2012 Q1	124	/	3293	3.77%	1	/	118	0.85%	78	/	2840	2.75%	207	/	6302	3.28%
2012 Q2	122	/	3298	3.70%	3	/	122	2.46%	96	/	2894	3.32%	223	/	6380	3.50%

Table 29.2 Proportion of individuals on antiretroviral therapy who change antiretroviral drug treatment by gender

Quarter	Male			Female		
	Count	Denominator	Percentage	Count	Denominator	Percentage
2006 Q1	279	3174	8.79%	44	525	8.38%
2006 Q2	351	3248	10.81%	55	543	10.13%
2006 Q3	349	3280	10.64%	67	557	12.03%
2006 Q4	406	3351	12.12%	52	569	9.14%
2007 Q1	346	3417	10.13%	62	589	10.53%
2007 Q2	327	3580	9.13%	61	607	10.05%
2007 Q3	216	3636	5.94%	63	627	10.05%
2007 Q4	325	3723	8.73%	68	628	10.83%
2008 Q1	309	3791	8.15%	57	650	8.77%
2008 Q2	387	3928	9.85%	71	683	10.40%
2008 Q3	280	4000	7.00%	62	724	8.56%
2008 Q4	259	4109	6.30%	57	732	7.79%
2009 Q1	260	4201	6.19%	64	774	8.27%
2009 Q2	202	4317	4.68%	67	803	8.34%
2009 Q3	178	4404	4.04%	48	804	5.97%
2009 Q4	195	4477	4.36%	32	832	3.85%
2010 Q1	194	4566	4.25%	40	854	4.68%
2010 Q2	169	4637	3.64%	49	871	5.63%
2010 Q3	198	4751	4.17%	52	893	5.82%
2010 Q4	188	4817	3.90%	58	926	6.26%
2011 Q1	191	4926	3.88%	64	953	6.72%
2011 Q2	162	4979	3.25%	57	988	5.77%
2011 Q3	195	5107	3.82%	52	998	5.21%
2011 Q4	150	5186	2.89%	50	1014	4.93%
2012 Q1	171	5269	3.25%	36	1033	3.48%
2012 Q2	175	5330	3.28%	48	1050	4.57%

Appendix A: List of Indicators

Indicator 1: Number of HIV tests

Indicator 2: Population HIV testing rate

Indicator 3: Number of new HIV diagnoses

Indicator 4: Rate of new AIDS case reports

Indicator 5: Percentage positivity among persons tested for HIV

Indicator 6a: Proportion of individuals tested for syphilis who are tested for HIV at the same clinical encounter

Indicator 6b: Proportion of individuals with a new STI diagnosis who are tested for HIV within three months of STI diagnosis

Indicator 7: Proportion of individuals with a new HCV diagnosis who are tested for HIV within three months of HCV diagnosis

Indicator 8: Proportion of individuals named as contacts of an index case who have a subsequent HIV test

Indicator 9: Proportion of individuals with a new HIV diagnosis with advanced HIV disease

Indicator 10: Proportion of HIV individuals with a CD4 cell count <200 cells/mL at first positive HIV test

Indicator 11: Proportion of individuals with a new HIV diagnosis with acute HIV infection

Indicator 12: Proportion of HIV positive individuals with a family physician

Indicator 13: Interval between first HIV positive test and first HIV plasma viral load

Indicator 14: Proportion of individuals starting antiretroviral therapy (ART) late in the course of HIV disease

Indicator 15: Proportion of HIV positive individuals eligible to start ART by CD4 cell count criteria who are currently on ART

Indicator 16: Proportion of HIV positive individuals who are accessing ART

Indicator 17: Rate of disease progression among individuals accessing and not accessing ART

Indicator 18: Proportion of individuals with a new HIV diagnosis who are tested for syphilis within three months of HIV diagnosis

Indicator 19: Proportion of HIV positive women with baseline and annual pap smears

Indicator 20: Proportion of HIV positive individuals who have had > three pVL measures in the past year

Indicator 21: Percentage of HIV-infected individuals who are tested for genotypic antiretroviral drug resistance prior to starting antiretroviral therapy (ART)

- Indicator 22:** Percentage of individuals starting antiretroviral therapy (ART) who achieve HIV plasma viral load (pVL) of <200 copies/mL within six months of therapy initiation
- Indicator 23:** Percentage of individuals who initiated antiretroviral therapy (ART) with a recommended therapy regimen (among those with no drug resistance)
- Indicator 24:** Percentage of individuals on antiretroviral therapy (ART) that achieve annual prescription refill adherence of >95%
- Indicator 25:** Number of physicians initiating antiretroviral therapy (ART)
- Indicator 26:** Percentage of individuals on antiretroviral therapy (ART) who experience a serious adverse drug reaction (ADR)
- Indicator 27:** Rate of transmission of primary resistance
- Indicator 28:** Incidence of resistance to any retroviral drug
- Indicator 29:** Proportion of individuals on antiretroviral therapy who change antiretroviral drug treatment
- Indicator 30:** Social determinant to be determined
- Indicator 31:** Social determinant to be determined
- Indicator 32:** Social determinant to be determined
- Indicator 33:** Social determinant to be determined
- Indicator 34:** Estimates of health service cost and use among HIV positive individuals using supportive services
- Indicator 35:** The number and spectrum of health care services used by HIV positive individuals who are using and not using ART
- Indicator 36:** Cost of health service use among HIV positive individuals who are using and not using ART